

PHYTOCHEMICAL ANALYSIS OF PETROLEUM ETHER EXTRACT OF GRATELOUPIA FILICINA (LAMOUROUX) C.AG. USING FTIR AND HPLC

¹Athi Lakshmi S, ²John Peter Paul J*.

¹*Research Scholar (Reg. No: 20211282262012), Department of Botany, St. Xavier's College (Autonomous), Palayamkottai-627 002, Tamil Nadu. Affiliated to Manonmanium Sundaranar University, Abishekapatti, Tirunelveli – 627 012, Tamil Nadu, India.

²Assistant Professor of Botany & Director, Centre for Advanced Research in Plant Sciences (CARPS), St. Xavier's College (Autonomous), Palayamkottai - 627 002, Tamil Nadu, India.

²Corresponding Author – johnarock2008@yahoo.com

Abstract:

Marine algae are a source of bioactive substances and they produce a great variety of secondary metabolites with a diverse range of biological activities. Phytochemical types include sterols, isoprenoids, amino acids, terpenoids, phlorotannins, steroids, phenolic compounds, halogenated ketones and alkanes, cyclic polysulphides, fatty acids and acrylic acid have been isolated from marine algae. Hence, pharmacologists, physiologists and chemists have been paying increasing attention to the marine organisms particularly on marine macro algae for screening bioactive substances. The present study was designed to discover the phytochemical analysis of petroleum ether extract of Grateloupia filicina (Lamouroux) C.Ag. collected from Manapad, Thoothukudi district, Tamil Nadu, India. The phytochemical screening of petroleum ether extract was carried out by using the standard procedure of FTIR spectroscopy and HPLC. The FTIR spectrum of the petroleum ether extract of Grateloupia filicina (Lamouroux) C.Ag. was revealed the presence of functional groups such as Aliphatic amines, Nitro compounds, Aromatics, Alkenes, Aldehydes, Esters, Alkanes and Carboxylic acids. The qualitative HPLC fingerprint profile displayed three compounds at different retention times of 1.293 min, 2.287min and 2.750 min indicate the presence of three compounds. The profile displayed the most prominent peak of maximum area percentage (93.4%) with the retention time of 2.287 min. The phytochemicals were characterized which showed the presence of secondary metabolites. The results of this study provide a base work for utilizing petroleum ether extract of Grateloupia filicina (Lamouroux) C.Ag., as a treatment for a variety of disorders.

Keywords: Grateloupia filicina, Macro algae, FTIR, HPLC.

Introduction

The Marine Ecosystem occupies one-third of the Earth's atmosphere which covers approximately 71 percent of the surface of the earth. In aquatic biodiversity Algae are categorised into microalgae and macro algae. A unique group of micro algae are blue-green algae, also called Cyanobacteria and the macro algae are classified as Chlorophyceae (green algae), Phaeophyceae (brown algae) and Rhodophyceae (red algae). It is classified based on the specific pigments. Marine algae are a source of bioactive substances, and they contain a huge variety of secondary metabolites with a wide range of biological activities. Over the past several decades seaweeds have been used by humans as medicine and food as a fresh source. Seaweeds are reservoirs of carotenoids, pigments, diverse functional polysaccharides. Seaweeds are excellent source of vitamin A, B1, B12, C, D and E (Skulberg, 2000). Seaweeds have generated an enormous amount of interest in the pharmaceutical industry as a source of bioactive compounds with immense medicinal potential (Shyamala and Thangaraju, 2013). The secondary metabolites synthesized by seaweeds demonstrate a broad spectrum of

bioactivity (Smith, 2004 and Craig, 1999). Due to the biological interest, the present study was undertaken to investigate the phytochemical analysis of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag., collected from Manapad, Thoothukudi district, Tamil Nadu, India.

MATERIALS AND METHODS

Collection of samples:

Grateloupia filicina (Lamouroux) C.Ag., the red algal species were collected from Manapad coastal region (Latitude: 8° 22' 36.5" N, Longitude: 78° 3' 29.968" E), Thoothukudi district, Tamil Nadu, India during the month of August 2021. The collected algal sample was authenticated by Dr. J. John Peter Paul, Assistant Professor of Botany & Director, Centre for Advanced Research in Plant Sciences (CARPS), St. Xavier's College (Autonomous), Palayamkottai and accumulated in Xavier's College Herbarium Centre for Biodiversity and Biotechnology, St. Xavier's College (Autonomous), Palayamkottai – 627002, and the given voucher number for the accumulated herbarium sample was XCH-20537. The samples were collected by hand picking during low waves and washed with marine water to remove debris and epiphytes. The entire epiphytes were turfed using a soft brush. In the laboratory, the samples were once again washed with freshwater followed by a rinse with distilled water and stored in a refrigerator for further analysis (Udhaya and Paul, 2017).

Preparation of petroleum ether extract

For the preparation of petroleum ether extract, the collected plant specimens were washed thoroughly and placed on blotting paper and spread out at room temperature in the shade condition for drying. The shade dried samples were grounded to fine powder using a tissue blender. The powdered samples were then stored in the refrigerator for further use. 30g powdered sample was packed in Soxhlet apparatus and extracted with petroleum ether for 8h separately. The excess amount of petroleum ether was evaporated and fine petroleum ether crude powder was prepared and stored in the refrigerator for the further analysis (John Peter Paul and Yuvaraj, 2013).

FTIR analysis

FTIR analysis of the petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag., was performed using Perkin Elmer Spectrophotometer system, which was used to detect the characteristic peaks and their functional groups. The peak values of the FTIR were recorded. Each and every analysis was repeated twice and confirmed the spectrum (John Peter Paul and Yuvaraj, 2013).

HPLC analysis

The HPLC analysis of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag., performed on a Shimadzu LC-10AT VP HPLC system, equipped with a model LC-10AT pump, UV-Visible detector SPD-10AT, a Rheodyne injector fitted with a 20µl loop and an auto injector SIL-10AT. A Hypersil® BDS C-18 column (4.6 × 250mm, 5µm size) with a C-18 guard column was used. The elution was carried out with gradient solvent systems with a flow rate of 1ml/min at ambient temperature (25-28°C). The mobile phase consisted of 0.1% v/v methanol (solvent A) and water (solvent B). The mobile phase was prepared daily, filtered through a 0.45µm and sonicated before use. Total running time was 15min. The sample

injection volume was 20 μ l while the wavelength of the UV-Visible detector was set at 254nm (John Peter Paul and Shri Devi, 2013).

Instrumentation

An isocratic HPLC (Shimadzu HPLC Class VP series) with two LC- 0 AT VP pumps (Shimadzu), a variable wave length programmable photo diode array detector SPD-M10A VP (Shimadzu), a CTO- 10AS VP column oven (Shimadzu), a SCL-10A VP system controller (Shimadzu), a reverse phase Luna 5 μ C18 (2) and Phenomenex column (250 mm X 4.6mm) were used. The mobile phase components ethanol:water (45:55) were filtered through a 0.2 μ membrane filter before use and were pumped from the solvent reservoir at a flow rate of 1ml/min which yielded column backup pressure of 260-270kgf/cm². The column temperature was maintained at 27°C. 20 μ l of the respective sample and was injected by using a Rheodyne syringe (Model 7202, Hamilton).

Result and Discussion

FTIR analysis

The FTIR spectrum was used to identify the functional group of the active components based on the peak value in the region of infra red radiation. The petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag. was passed into the FTIR and the functional groups of the components were separated based on its peak ratio. The FTIR results of petroleum ether extract of *Grateloupia filicina* showed different peaks at 1081.99, 1338.51, 1489.91, 1680.85, 1724.24, 1768.60, 2820.70, 2883.38 and 2944.13 cm⁻¹. the functional groups such as aliphatic amines, nitro compounds, aromatics, alkenes, aldehydes, esters, aldehydes, alkanes and carboxylic acids respectively (Figure 1& Table 1).

Table-1: FTIR peak values and functional groups of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag.

S.No.	Frequency (cm ⁻¹)	Functional group	Possible compounds
1.	1081.99	C–N	Aliphatic amines
2.	1338.51	N–O	Nitro compounds
3.	1489.91	C–C	Aromatics
4.	1680.85	–C=C–	Alkenes
5.	1724.24	C=O	Aldehydes
6.	1768.60	C=O	Esters
7.	2820.70	H–C=O	Aldehydes
8.	2883.38	C–H	Alkanes
9.	2944.13	O–H	Carboxylic acids




Figure-1: FTIR spectrum of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag.

HPLC analysis

The qualitative HPLC fingerprint profile of the petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag. was prepared by hot extraction was subjected to HPLC for the separation and identification of constituents present in the *Grateloupia filicina*. Three compounds were separated at different retention time of 1.293 min, 2.287min and 2.750 min respectively. The profile displayed one prominent peak maximum area percentage (93.4%) with the retention time of 2.287 min (Figure 2& Table 2).

Table-2: HPLC profile of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag.

S.No.	Retention time (min)	Area (Mv.s)	Height (mV)	Area (%)	Height (%)	W05(min)
1.	1.293	0.422	0.082	1.2	1.2	0.08
2.	2.287	32.122	6.208	93.4	93.4	0.08
3.	2.750	1.839	0.353	5.3	5.3	0.08
	Total	34.383	6.643	100.0	100.0	




Figure-2: HPLC spectrum of petroleum ether extract of *Grateloupia filicina* (Lamouroux) C.Ag.

CONCLUSION

From the present study, it was concluded that, FTIR and HPLC analysis can be used as effective tool in identifying the phytochemicals. It also suggested that *Grateloupia filicina* is the richest sources of phytochemicals which can be isolated and further screened for different kinds of biological activities depending on the therapeutic uses. Further work will be conducted the isolation and characterization of active principles responsible for the biopotential. The presence of various functional groups and phytocompounds in *Grateloupia filicina* conform that it acts as a most important source of drugs against various ailments.

REFERENCE

1. Skulberg, O.M. Microalgae as a source of bioactive molecules-experience from cyanophyte research. *Journal of Applied Phycology*. 2000, 12(3-5):341-348.
2. Shyamala, V., Thangaraju, N. Screening of phytochemical and antibacterial activity of three different seaweeds from Gulf of Mannar, Tamil Nadu. *Phykos*, 2013, 43(1):32-38.
3. Smith, A.J. Medicinal and pharmaceutical uses of seaweed natural products: A review. *Journal of Applied Phycology*, 2004, 16:245-262.
4. Craig, W.J. Health-promoting properties of common herbs. *Am J Clin Nutr.* 1999; 70:491S-499S.
5. Udhaya C Iniya, Paul J John Peter. 2017. Screening of preliminary phytochemicals of *Gracilaria cylindrica* Boergesen in Koothankuzhi, Tirunelveli district, Tamil Nadu, India. *Indo American Journal of Pharmaceutical Sciences*,; 4(12):4590-4594.
6. John Peter Paul J, Yuvaraj P. (2013). Phytochemical analysis of *Padina distromatica* Hauck. *Indo American Journal of Pharmaceutical Research*, 3(7), 5290-5297.
7. John Peter Paul, J., Shri Devi, S.D.K. Phytochemical studies on *Padina pavonica* (L.) Thivy ex Taylor. *Indo American Journal of Pharmaceutical Research*, 2013, 3(6):4452-4459.