Ijfans International Journal of Food and Nutritional Sciences
ISSN PRINT 2319 1775 Online 2320 7876
Research Paper © 2012 IJFANS. All Rights Reserved Journal UGC CARE Listed (Group-I) Volume 10, Issue 12, Dec 2021

Green Synthesis of Biologically Active Organic Compounds Using Eco-Friendly Catalysts: A Comprehensive Review

Shinde Pooja Bhagirath

Assistant Professor in Department of Chemistry, Nutan vidya prasarak Mandal's Arts, Commerce and Science College Lasalgaon, Taluka Niphad, District Nashik, Maharashtra, India 422306

Abstract

The synthesis of biologically active organic compounds has traditionally relied on hazardous solvents, stoichiometric reagents, and energy-intensive processes that generate substantial chemical waste. This review examines the paradigm shift toward environmentally sustainable methodologies, focusing on eco-friendly catalysts that enable the efficient construction of bioactive molecular frameworks while minimizing environmental impact. Drawing upon recent advances from 2010–2020, the paper synthesizes evidence across catalyst classes including recyclable Brønsted acid systems, heterogeneous nanocatalysts, deep eutectic solvents (DES), supported ionic liquids, and photo/electrocatalytic systems. Notable achievements include caffeine–H_3PO_4 as a recyclable metal-free catalyst for fused pyrimidines with demonstrated antibacterial and cytotoxic activity ; graphene oxide/ZnO nanocomposites enabling solvent-free Biginelli synthesis of dihydropyrimidinones with 69–97% yields and four-cycle recyclability ; benzethonium chloride:glycerol DES achieving 77–93% yields of benzimidazole-fused pyrimidines with anti-diabetic IC_{50} values of 25–44 µg/mL and five-cycle reuse ; and Merrifield resin-supported ionic liquids affording ≥97% yields of xanthenediones within ≤5 minutes under ultrasound at ambient temperature, with eight-cycle recyclability . The review evaluates green metrics including Atom Economy, E-Factor, and Process Mass Intensity, demonstrating that these methodologies can achieve E-factors below 1, outperforming typical laboratory-scale syntheses . Critical gaps remain in the scalability of nanocatalyst preparation, long-term catalyst stability under industrial conditions, and the integration of life cycle assessment into early-stage method development. The paper concludes that eco-friendly catalysis has matured from proof-of-concept demonstrations to a robust toolkit for sustainable bioactive compound synthesis, though translation to industrial practice requires further validation.

Keywords:

Green synthesis, eco-friendly catalysts, biologically active compounds, deep eutectic solvents, nanocatalysts, recyclable catalysts, green metrics, sustainable organic synthesis

1. Introduction

The pharmaceutical and fine chemical industries stand at a critical juncture. While they deliver products essential to human health and well-being, their environmental footprint is substantial: the pharmaceutical sector generates approximately 49 tons of CO_2 per million dollars of revenue, surpassing even automotive manufacturing in emissions intensity . Traditional organic

Research Paper © 2012 IJFANS. All Rights Reserved

synthesis—the engine of drug discovery and development—relies heavily on volatile organic solvents, stoichiometric activating agents, and energy-intensive heating, producing large volumes of hazardous waste relative to product mass.

Green chemistry offers a systematic framework for addressing these challenges. The twelve principles articulated by Anastas and Warner provide a design philosophy: prevent waste, maximize atom economy, use safer solvents and auxiliaries, design for energy efficiency, and employ catalytic rather than stoichiometric reagents, among others . Within this framework, catalysis occupies a privileged position. Catalysts reduce the quantity of reagents required, often enable milder reaction conditions, and can be recovered and reused—directly addressing the waste and energy challenges that characterize conventional synthesis.

This review examines the application of eco-friendly catalysts to the synthesis of biologically active organic compounds, with emphasis on methods that demonstrate measurable environmental benefits alongside biological efficacy. The scope encompasses heterogeneous nanocatalysts, deep eutectic solvents, supported ionic liquids, recyclable Brønsted acid systems, and unconventional activation methods (microwave, ultrasound, photoredox) that complement catalytic approaches. The objective is to assess the state of the art, identify the most promising catalyst systems, evaluate their green credentials through quantitative metrics, and delineate the barriers to broader adoption.

2. Definitions

Green Synthesis: The design of chemical products and processes that reduce or eliminate the use or generation of hazardous substances, guided by the twelve principles of Green Chemistry. In the context of bioactive compound synthesis, green synthesis emphasizes reduced solvent consumption, recyclable catalysts, energy-efficient activation, and minimal waste generation.

Eco-Friendly Catalyst: A catalytic material or system that facilitates a chemical transformation while minimizing environmental burden. Criteria include low toxicity, recoverability and reusability, high selectivity, and avoidance of precious or scarce elements. Examples include metal-free organic catalysts, biodegradable deep eutectic solvents, and recyclable heterogeneous nanocomposites.

Green Metrics: Quantitative measures of the environmental performance of a synthetic process. Key metrics include:

1. **Atom Economy (AE):** Molecular weight of desired product divided by total molecular weight of all reactants, expressed as a percentage.
2. **E-Factor:** Mass of waste divided by mass of product. Lower values indicate greater efficiency; E-factors below 1 are considered excellent for fine chemicals .
3. **Process Mass Intensity (PMI):** Total mass of materials used divided by mass of product.
4. **EcoScale:** A composite metric penalizing hazards, energy use, and waste.

Deep Eutectic Solvent (DES): A mixture of two or more components—typically a hydrogen bond acceptor (HBA) and a hydrogen bond donor (HBD)—that forms a liquid with a melting point significantly lower than that of its constituents. DESs serve as both reaction media and

Research Paper © 2012 IJFANS. All Rights Reserved

catalysts, offering non-volatile, non-flammable, and often biodegradable alternatives to conventional organic solvents .

Recyclability: The capacity of a catalyst to be recovered after a reaction and reused for subsequent cycles without significant loss of activity or selectivity. Recyclability is a key determinant of the environmental and economic viability of catalytic processes.

3. Need for the Study

Three considerations establish the imperative for this review.

First, the environmental burden of conventional bioactive compound synthesis is unsustainable. The pharmaceutical industry's waste intensity—measured by E-factors often exceeding 25 for fine chemicals and 100 for pharmaceuticals—contrasts sharply with the commodity chemical sector, where E-factors are typically below 5. The development of synthetic methods that achieve comparable yields and selectivity with dramatically reduced waste is therefore a priority for both environmental and economic reasons.

Second, the catalyst landscape has diversified significantly in recent years, yet comparative assessments across catalyst classes are scarce. Researchers can choose from metal-free organic acids, heterogeneous metal oxides, supported ionic liquids, DESs, and photo/electrocatalytic systems, but the evidence base for selecting among these options—based on green metrics, biological activity of products, and practical considerations—is fragmented.

Third, the translation of green synthetic methods from laboratory demonstration to industrial practice requires more than high yields. Catalyst stability, scalability, cost, and life cycle impacts must be evaluated. This review synthesizes the available evidence on these dimensions to identify both the most promising catalyst systems and the critical gaps that remain.

4. Aims and Objectives

This review aims to provide a comprehensive, evidence-based assessment of eco-friendly catalysts for the synthesis of biologically active organic compounds, evaluating their synthetic performance, green credentials, and the biological activities of the products they enable.

Objectives:

1. To categorize and evaluate the major classes of eco-friendly catalysts employed in bioactive compound synthesis.
2. To quantify synthetic performance (yields, reaction times, recyclability) and green metrics (E-factor, atom economy) for representative catalyst systems.
3. To document the biological activities (antimicrobial, anticancer, antidiabetic, etc.) of products synthesized using these methods.
4. To identify the advantages and limitations of each catalyst class.
5. To assess the barriers to industrial adoption and propose a research agenda for translation.

5. Hypothesis

Research Paper © 2012 IJFANS. All Rights Reserved

H1: Eco-friendly catalysts can achieve synthetic performance (yields, selectivity, reaction times) comparable to or exceeding conventional methods while substantially reducing waste generation and hazardous reagent use.

H2: Catalyst recyclability is the single most important determinant of the environmental and economic viability of green synthetic methods for bioactive compounds.

H3: The biological activities of compounds synthesized using eco-friendly catalysts are not compromised relative to those produced by conventional methods; in some cases, green methods may enable access to compounds not accessible via traditional routes.

6. Literature Search

The literature on eco-friendly catalysis for bioactive compound synthesis spans heterogeneous catalysis, green chemistry, medicinal chemistry, and materials science. This review synthesizes key contributions from 2010–2020.

Recyclable Brønsted Acid Catalysts. A caffeine–H_3PO_4 system has been developed as a recyclable, metal-free acidic catalyst for the synthesis of pyrimido[4,5-d]pyrimidine derivatives under ethanolic conditions . The catalyst was characterized by FT-IR, NMR, XRD, SEM, EDX, TGA, DTG, DTA, and DSC, confirming stability and robustness. It exhibited excellent recyclability across multiple runs with negligible activity loss. Biological screening demonstrated significant cytotoxic potential, supported by molecular docking, pharmacokinetic, and ADMET analyses . Mechanistic studies indicate cooperative dual-functionality: phosphoric acid activates the aldehyde carbonyl through protonation, while the caffeine scaffold stabilizes reactive intermediates via π–π stacking and hydrogen bonding, lowering activation barriers for Knoevenagel condensation and subsequent cyclization .

Heterogeneous Nanocatalysts. Graphene oxide/ZnO nanocomposites (GO/ZnO-NC) have been employed as heterogeneous catalysts for the solvent-free Biginelli synthesis of dihydropyrimidinones/thiones (DHPMs/DHPMTs) . ZnO nanoparticles were green-synthesized using sugarcane stem extract, and the GO/ZnO nanocomposite provided higher yields than GO or ZnO alone, achieving 69–97% yields with four-cycle recyclability. The method represents a solvent-free, low-cost, energy-efficient approach to bioactive heterocyclic products . A review in *Organic & Biomolecular Chemistry* documents the broader use of nanostructured catalysts for constructing drug-like hybrid molecules with privileged heterocyclic substructures, emphasizing their role in accessing natural product mimetic scaffolds under sustainable conditions .

Deep Eutectic Solvents. Benzethonium chloride:glycerol (1:2) DES has been employed as both reaction medium and catalyst for the one-pot multicomponent synthesis of ethyl 2-methyl-4-aryl-1,4-dihydrobenzo[4,5]imidazo[1,2-a]pyrimidine-3-carboxylate derivatives . The reaction proceeded at 110°C, affording 11 derivatives in 77–93% yields within 45–90 minutes. The DES was recovered by water addition and reused across five consecutive cycles with sustained catalytic activity. Products exhibited dose-dependent α-amylase inhibitory activity with IC_{50} values of 25–44 µg/mL, indicating anti-diabetic potential . A Brønsted-based DES comprising betaine and p-toluenesulfonic acid ($[Bet]_2[PTSA]$) has been applied to the synthesis

Research Paper © 2012 IJFANS. All Rights Reserved

of 1,2,4,5-tetrasubstituted imidazoles with cytotoxic activity against HepG2 liver carcinoma cells . The DES serves as a non-volatile, recyclable Brønsted acid catalyst, eliminating the need for external catalysts and corrosive liquid acids.

Supported Ionic Liquids. N-methylbenzimidazolium-functionalized Merrifield resin-supported ionic liquids ([MRBIm][X]) have been evaluated as recyclable heterogeneous organocatalysts for the ultrasound-assisted synthesis of 1,8-dioxo-octahydroxanthenes . The [MRBIm][HSO_4] catalyst demonstrated superior activity, affording open-chain products in ≥97% yields within ≤5 minutes at 25°C. At 60°C, the reaction pathway shifted toward closed-chain derivatives via intramolecular aldol cyclization, demonstrating temperature-dependent selectivity. The catalyst was recovered and reused for eight runs with consistent performance .

Recyclable Metal Oxide Catalysts. Commercial MgO has been employed as a recyclable heterogeneous catalyst for the microwave-assisted, solvent-free synthesis of antiviral furano-chalcones from biomass-derived furanic aldehydes . The catalyst maintained activity and selectivity over six reuse cycles. Green metrics (Atom Economy, PMI, E-factor, EcoScale, Green Motion™) demonstrated superior performance compared to conventional solvent-based methods, with E-factor <1 attributable to solvent elimination, recyclable catalysis, and microwave efficiency .

Unconventional Activation Methods. Visible-light photoredox catalysis enables radical-mediated transformations under mild conditions, with applications in benzopyran (coumarin, flavone, chromone, chromene, chromane) synthesis . Microwave irradiation provides homogeneous heating with reduced reaction times. Sonochemistry (ultrasound) accelerates reactions via cavitation. Mechanochemistry enables solvent-free transformations through direct mechanical energy transfer. Continuous-flow chemistry improves parameter control and safety. Electrochemistry uses electrons as reactants, eliminating stoichiometric oxidants or reductants . These methods complement catalytic approaches, often enabling further reductions in energy consumption and waste.

Green Metrics and Life Cycle Considerations. A comprehensive whitepaper on green chemistry in organic synthesis details the application of mass-based metrics (Atom Economy, E-Factor, PMI) and advanced frameworks (WAR Algorithm, GREENSCOPE) for evaluating environmental and economic viability . The importance of Life Cycle Assessment (LCA) is emphasized to ensure that improvements in synthetic efficiency do not cause upstream or downstream ecological harm. Green Analytical Chemistry tools (NEMI, GAPI, AGREE) are also discussed for assessing the ecological footprint of analytical procedures .

7. Research Methodology

This review employs a **systematic narrative synthesis** of peer-reviewed studies published primarily between 2010 and 2020, focusing on eco-friendly catalytic methods for the synthesis of biologically active organic compounds.

Sources. The analysis draws upon: (1) studies in *Wiley Online Library*, *RSC Publishing*, *ACS Publications*, *ScienceDirect*, and *Europe PMC*; (2) catalyst systems including caffeine–H_3PO_4, GO/ZnO nanocomposites, benzethonium chloride:glycerol DES, $[Bet]_2[PTSA]$ DES,

Research Paper © 2012 IJFANS. All Rights Reserved

Merrifield resin-supported ionic liquids, and MgO; (3) unconventional activation methods (photoredox, microwave, ultrasound, mechanochemistry) as documented in the benzopyran review ; and (4) green metrics frameworks and LCA considerations .

Analytical Framework. The review is organized by catalyst class. For each system, the following dimensions are assessed: (a) synthetic performance (yield, reaction time, substrate scope); (b) recyclability (number of cycles, activity retention); (c) green metrics (where reported); (d) biological activity of products; and (e) practical limitations.

Limitations. The review is constrained by the heterogeneity of green metric reporting across studies. Many papers report yields and recyclability but not E-factor or PMI. Biological activity data vary in assay type and rigor. The review does not conduct original experimental work or meta-analysis.

8. Strong Points: Demonstrated Advantages of Eco-Friendly Catalysts

Comparable or Superior Synthetic Performance. The catalyst systems reviewed achieve yields that rival or exceed conventional methods. Merrifield resin-supported ionic liquids afford ≥97% yields within ≤5 minutes . GO/ZnO nanocomposites achieve 69–97% yields in solvent-free Biginelli reactions . Benzethonium chloride:glycerol DES delivers 77–93% yields within 45–90 minutes . These performance metrics demonstrate that green methods do not require sacrificing efficiency.

Excellent Recyclability. Recyclability is a consistent strength across catalyst classes. Merrifield-supported ionic liquids maintain performance over eight cycles . MgO retains activity over six cycles . Benzethonium chloride:glycerol DES and caffeine–H_3PO_4 each demonstrate five-cycle reuse with negligible activity loss . GO/ZnO nanocomposites are reusable over four cycles . This recyclability directly reduces catalyst consumption and waste generation.

Demonstrated Biological Activity of Products. The bioactive compounds synthesized using these methods exhibit meaningful biological activities. Fused pyrimidines from caffeine–H_3PO_4 catalysis show cytotoxic potential with favorable ADMET profiles . Benzimidazole-fused pyrimidines from DES catalysis exhibit α-amylase inhibition with IC_{50} values of 25–44 μg/mL, indicating anti-diabetic potential . Tetrasubstituted imidazoles from $[Bet]_2[PTSA]$ DES show cytotoxicity against HepG2 cells . Furano-chalcones from MgO catalysis are designed as antiviral agents . These findings confirm that green synthesis does not compromise biological efficacy.

Quantified Environmental Benefits. The MgO-catalyzed furano-chalcone synthesis reports an E-factor below 1, outperforming typical laboratory-scale syntheses . The combination of solvent elimination, recyclable heterogeneous catalysis, and microwave heating achieves waste minimization that is quantifiably superior to conventional solvent-based methods. The green metrics framework provided by the Zenodo whitepaper enables systematic comparison and identification of improvement opportunities .

Research Paper © 2012 IJFANS. All Rights Reserved

Access to Privileged Scaffolds. The methods reviewed enable access to medicinally privileged heterocyclic scaffolds including pyrimidines, benzopyrans, xanthenes, and chalcones . Nanocatalyzed syntheses facilitate the construction of hybrid heterocyclic scaffolds with structural diversity and molecular complexity, supporting drug discovery efforts . The ability to generate natural product mimetic scaffolds under sustainable conditions expands the accessible chemical space for bioactive compound discovery.

Methodological Diversity. The field offers multiple catalyst options, enabling selection based on substrate, scale, and available equipment. Metal-free organic acids (caffeine–H_3PO_4) avoid metal contamination concerns . Magnetic nanocatalysts offer facile recovery . DESs provide dual solvent-catalyst function with biodegradability . Supported ionic liquids combine high activity with heterogeneous recovery . This diversity allows researchers to match catalyst choice to specific requirements.

9. Weak Points: Limitations and Barriers

Scalability of Catalyst Preparation. Many nanocatalyst systems require multistep preparation involving specialized equipment and reagents. GO/ZnO nanocomposites involve Hummer's method for GO synthesis followed by green ZnO nanoparticle preparation and composite formation . Merrifield resin-supported ionic liquids require functionalization of the polymer support . These preparation steps add cost and complexity that may limit adoption at scale. The ACS review notes that nanocatalyst-mediated strategies, while offering optimal balance of yield and recyclability, involve "multistep synthesis" as a limitation .

Limited Green Metric Reporting. While some studies report E-factor or PMI, many do not . The absence of standardized green metric reporting hinders comparative assessment across catalyst classes. The Zenodo whitepaper advocates for systematic use of mass-based metrics and frameworks including WAR Algorithm and GREENSCOPE, but adoption in primary research remains inconsistent .

Long-Term Stability and Leaching. Studies typically report recyclability over 4–8 cycles, but the long-term stability of catalysts under repeated use—particularly metal leaching from supported systems—is less thoroughly characterized. For pharmaceutical applications, even trace metal contamination can be problematic, requiring rigorous quality control. Metal-free catalysts (caffeine–H_3PO_4, DESs) offer advantages in this regard but may have different stability profiles.

Substrate Scope Limitations. Many studies demonstrate scope across 10–15 derivatives, but broader substrate tolerance—particularly for functionally diverse or sterically hindered substrates—may be limited. The ACS review notes that solvent-free methods show "slight yield reduction with ortho-substituted and sterically hindered substrates" . Expanding substrate scope while maintaining green credentials is an ongoing challenge.

Energy Input Considerations. While microwave and ultrasound methods reduce reaction times, they require specialized equipment with energy demands that may offset some green benefits depending on the energy source. The green credentials of these methods are contingent

Research Paper © 2012 IJFANS. All Rights Reserved

on the carbon intensity of electricity supply. Life cycle assessment, as advocated in the Zenodo whitepaper, is necessary to evaluate net environmental impact .

Translation Gap to Industrial Practice. The reviewed studies are predominantly laboratory-scale (millimole to gram quantities). Translation to kilogram or ton scale introduces challenges in mixing, heat transfer, catalyst recovery, and process safety that are not addressed in small-scale demonstrations. The ACS review acknowledges that for ultrasound-assisted methods, "scalability from laboratory to industrial scale remains challenging" .

Biological Evaluation Depth. While products demonstrate biological activity in preliminary assays, most studies report only initial screening (cytotoxicity, antimicrobial, enzyme inhibition). Mechanistic studies, structure-activity relationship analysis, and in vivo validation are less common. The caffeine–H_3PO_4 study includes molecular docking and ADMET analysis , but such comprehensive characterization is not universal.

10. Current Trends

Deep Eutectic Solvents as Dual-Function Systems. DESs are increasingly employed as both reaction media and catalysts, eliminating the need for separate solvents and catalysts while simplifying workup and enabling recyclability . The betaine:PTSA system demonstrates that DES acidity can be tuned by HBD selection, enabling Brønsted acid catalysis without corrosive liquid acids .

Magnetic Nanocatalysts for Facile Recovery. Fe_3O_4-based magnetic nanocatalysts offer recovery by external magnet, simplifying separation and reducing catalyst loss . Immobilized Schiff base–Mn complexes on magnetic supports have been applied to the synthesis of biologically active pyrido[1,2-a]pyrimidin-6-ones in aqueous media .

Solvent-Free and Minimal-Solvent Conditions. Mechanochemistry, solvent-free Biginelli reactions, and solventless microwave synthesis reduce or eliminate organic solvent use . The catalyst-free, microwave-assisted synthesis of benzylidene Schiff bases under solventless conditions demonstrates that even catalyst-free methods can achieve green objectives when combined with efficient activation .

Photo and Electrochemical Activation. Visible-light photoredox catalysis and electrochemistry use photons or electrons as clean reagents, eliminating stoichiometric oxidants or reductants . These methods are particularly attractive for radical-mediated transformations that are difficult to achieve under conventional thermal conditions.

Integration of Computational and Experimental Approaches. Molecular docking, pharmacokinetic prediction, and ADMET analysis are increasingly integrated into green synthesis studies, enabling early assessment of biological potential and guiding structure optimization . This integration accelerates the identification of promising candidates from green synthetic routes.

Emphasis on Green Metrics and LCA. There is growing recognition that yield alone is an insufficient measure of sustainability. The adoption of E-factor, PMI, and LCA frameworks is

Research Paper © 2012 IJFANS. All Rights Reserved

increasing, though unevenly . The RSC paper on MgO-catalyzed furano-chalcones explicitly benchmarks green metrics against conventional methods, demonstrating E-factor <1 .

11. History

The application of eco-friendly catalysts to bioactive compound synthesis has evolved through several phases, reflecting broader developments in green chemistry and catalysis.

Early Green Chemistry Foundations (1990s–2000s). The articulation of the twelve principles of green chemistry provided the conceptual framework for sustainable synthesis. Early applications focused on solvent replacement (water, ionic liquids) and the use of recyclable solid acids. Catalysis was recognized as a foundational green strategy, but practical demonstrations for complex bioactive molecules were limited.

Ionic Liquids and Supported Catalysts (2000s–2010s). Ionic liquids emerged as tunable, non-volatile solvents with catalytic potential. Supported ionic liquids addressed recovery and reuse challenges. Heterogeneous catalysts—zeolites, clays, metal oxides—were applied to multicomponent reactions relevant to heterocyclic synthesis. However, concerns about ionic liquid toxicity and cost prompted interest in alternative media.

Deep Eutectic Solvents and Nanocatalysts (2010s–2020s). DESs emerged as low-cost, biodegradable alternatives to ionic liquids, with the ability to serve as both solvent and catalyst. Nanocatalysts—particularly magnetic nanoparticles and graphene-based composites—offered high surface area, facile recovery, and tunable activity. The convergence of these materials with multicomponent reactions enabled rapid access to diverse heterocyclic libraries under sustainable conditions.

Current Era (2010–2020): Metrics, Mechanistic Understanding, and Biological Validation. The most recent phase is characterized by three developments. **First**, green metrics are being systematically applied, moving beyond qualitative claims of "greenness" to quantitative assessment . **Second**, mechanistic understanding of catalyst function—including the cooperative dual-functionality of caffeine–H_3PO_4 and the hydrogen-bonding activation by DESs—is enabling rational catalyst design . **Third**, biological validation is increasingly integrated into method development, ensuring that green synthetic routes deliver compounds with genuine therapeutic potential.

12. Discussion

The evidence assembled in this review supports a clear conclusion: **eco-friendly catalysis has matured into a robust and versatile toolkit for the synthesis of biologically active organic compounds, achieving synthetic performance comparable to conventional methods while delivering quantifiable environmental benefits**. However, translation to industrial practice and systematic integration of green metrics remain incomplete.

The Recyclability Advantage. Across catalyst classes, recyclability emerges as the single most important determinant of environmental and economic viability. The ability to recover and reuse catalysts over 5–8 cycles dramatically reduces catalyst consumption, waste generation, and cost. Merrifield-supported ionic liquids maintaining performance over eight

Research Paper © 2012 IJFANS. All Rights Reserved

cycles , MgO over six cycles , and DESs over five cycles demonstrate that recyclability is not an aspiration but a demonstrated characteristic. For industrial adoption, however, longer-term stability data (20+ cycles) and mechanistic understanding of deactivation pathways are needed.

Green Metrics as Decision Tools. The E-factor below 1 achieved in MgO-catalyzed furano-chalcone synthesis provides a benchmark that should inform method selection. Conventional bioactive compound syntheses often generate 10–100 times more waste per unit product. The systematic application of E-factor, PMI, and LCA frameworks enables researchers to quantify the benefits of green methods and identify hotspots for improvement. The integration of metrics into early-stage method development—rather than as a post-hoc assessment—would accelerate progress.

The Metal-Free Advantage. Metal-free catalysts—caffeine–H_3PO_4, DESs, organic acids—offer particular advantages for pharmaceutical applications where metal contamination is tightly regulated. The caffeine–H_3PO_4 system achieves this while maintaining excellent recyclability and demonstrating biological activity in products . The dual-functionality mechanism—acid catalysis combined with molecular stabilization by caffeine—illustrates how rational design can achieve performance without metals .

Biological Validation as a Non-Negotiable Requirement. The ultimate purpose of synthesizing bioactive compounds is to discover and develop therapeutic agents. The studies reviewed demonstrate that green methods yield compounds with genuine biological activity: anti-diabetic DES-derived pyrimidines , cytotoxic fused pyrimidines , anticancer imidazoles , and antiviral furano-chalcones . The integration of computational (docking, ADMET) and experimental (in vitro assays) validation ensures that green synthesis is not merely an academic exercise but a pathway to real therapeutic candidates.

Remaining Barriers. Three barriers stand between laboratory demonstration and industrial adoption. **Scalability** of nanocatalyst preparation and heterogeneous catalytic processes requires engineering solutions that are not addressed in millimole-scale studies. **Regulatory acceptance** of green processes and catalysts may require validation against established methods, particularly for pharmaceutical manufacturing. **Economic competitiveness** depends on catalyst cost, recyclability, and the value of waste reduction—factors that vary by scale and application. The ACS review's comparative table provides a useful starting point for technology selection, identifying nanocatalyst-mediated and microwave-assisted solvent-free protocols as the most sustainable and efficient approaches, while acknowledging that classical methods remain relevant for large-scale cost-sensitive applications .

13. Results

Recyclable Brønsted Acid (Caffeine–H_3PO_4). Fused pyrimidines synthesized in notable yields within short reaction times. Catalyst stable and recyclable across multiple runs. Products show significant cytotoxic potential with favorable drug-like attributes in ADMET analysis . Docking scores against CDK2 ranged from −8.7 to −10.8 kcal/mol .

Research Paper © 2012 IJFANS. All Rights Reserved

Heterogeneous Nanocatalyst (GO/ZnO). Solvent-free Biginelli synthesis of DHPMs/DHPMTs: 69–97% yields, four-cycle recyclability. ZnO nanoparticles green-synthesized from sugarcane stem extract. Higher yields than GO or ZnO alone .

Deep Eutectic Solvents. Benzethonium chloride:glycerol DES: 77–93% yields, 45–90 minutes, five-cycle reuse, anti-diabetic IC_{50} 25–44 μg/mL . $[Bet]_2[PTSA]$ DES: tetrasubstituted imidazoles with cytotoxicity against HepG2 cells .

Supported Ionic Liquids. [MRBIm][HSO_4]: ≥97% yields of xanthenediones within ≤5 minutes at 25°C. Temperature-dependent selectivity (open-chain at 25°C, closed-chain at 60°C). Eight-cycle recyclability .

Recyclable Metal Oxide. MgO: furano-chalcones from biomass-derived aldehydes, six-cycle reuse, E-factor <1, superior green metrics (AE, PMI, EcoScale) vs. conventional methods .

Unconventional Activation. Photoredox, microwave, ultrasound, mechanochemistry, continuous flow, and electrochemistry each offer specific advantages for benzopyran synthesis, enabling milder conditions and reduced waste .

Comparative Assessment. Nanocatalyst-mediated and microwave-assisted solvent-free protocols identified as most sustainable and efficient. Classical methods remain relevant for cost-sensitive large-scale applications .

14. Conclusion

The synthesis of biologically active organic compounds using eco-friendly catalysts has transitioned from a niche academic interest to a mainstream research enterprise. The evidence assembled in this review demonstrates that **catalyst systems including recyclable Brønsted acids, heterogeneous nanocomposites, deep eutectic solvents, supported ionic liquids, and recyclable metal oxides can achieve yields, selectivity, and reaction times comparable to or exceeding conventional methods while delivering quantifiable reductions in waste and hazardous reagent use**.

The most compelling findings are the demonstrated recyclability across multiple cycles—eight for supported ionic liquids , six for MgO , five for DESs —and the biological activity of the products, which validates that green methods access genuine therapeutic candidates rather than mere chemical curiosities. The E-factor below 1 achieved in MgO-catalyzed synthesis establishes a benchmark for what is achievable when solvent elimination, recyclable catalysis, and efficient activation are combined.

Yet the field faces critical gaps. Green metric reporting remains inconsistent, hindering comparative assessment. Long-term catalyst stability beyond 8 cycles is largely unexplored. Scalability from millimole to industrial scale requires engineering solutions that laboratory studies do not address. Life cycle assessment, while advocated , is rarely integrated into primary research. And the regulatory and economic frameworks for adopting green processes in pharmaceutical manufacturing remain underdeveloped.

The path forward requires coordinated effort: standardized green metric reporting as a condition for publication; systematic stability studies under industrially relevant conditions;

Research Paper © 2012 IJFANS. All Rights Reserved

engineering research on catalyst recovery and process intensification; and collaboration between academic method developers and industrial process chemists. The potential rewards—pharmaceutical manufacturing that is both economically competitive and environmentally sustainable—are substantial enough to justify the investment. The catalysts exist; the methods work; the products are bioactive. What remains is the translation.

15. Suggestions and Recommendations

For Researchers:

1. **Report green metrics routinely.** E-factor, PMI, and Atom Economy should accompany yield and recyclability data in all publications on green synthesis. The frameworks are well-established ; the barrier is adoption, not availability.
2. **Extend recyclability studies beyond 8 cycles.** Industrial application requires catalysts that maintain activity over dozens or hundreds of cycles. Studies that stop at 4–8 cycles leave a critical gap in the evidence base.
3. **Integrate LCA from the outset.** Rather than assessing environmental impact post-hoc, LCA should inform catalyst selection, solvent choice, and process design from the beginning .
4. **Validate biological activity mechanistically.** Preliminary screening is necessary but insufficient. Structure-activity relationship studies, target identification, and in vivo validation are needed to establish genuine therapeutic potential.

For Industrial Process Chemists:

5. **Evaluate nanocatalyst scalability.** The preparation of GO/ZnO, magnetic nanocatalysts, and supported ionic liquids requires multistep synthesis. Process research should focus on simplified preparation, quality control, and cost reduction.
6. **Consider DESs for Brønsted acid catalysis.** The tunability, biodegradability, and dual solvent-catalyst function of DESs offer practical advantages for reactions requiring acid catalysis, with demonstrated recyclability .

For Funding Agencies and Policymakers:

7. **Support translation research.** Grants that bridge laboratory method development and industrial process validation are needed to realize the practical benefits of green catalysis.
8. **Develop green chemistry education and training.** The principles and metrics of green chemistry should be integrated into graduate curricula in organic synthesis, catalysis, and process chemistry.

16. Future Scope

1. **Continuous-flow green synthesis.** Translation of batch recyclable catalyst systems to continuous-flow processes could improve productivity, safety, and catalyst lifetime.

Research Paper © 2012 IJFANS. All Rights Reserved

2. **Photo and electrocatalytic scale-up.** The mild conditions of photoredox and electrochemical activation are attractive, but reactor design and scale-up remain challenging. Engineering research is needed.
3. **Machine learning for catalyst selection.** The growing body of green synthesis data could support data-driven prediction of optimal catalyst-substrate combinations, reducing experimental trial and error.
4. **Biocatalysis integration.** Enzymes offer exquisite selectivity under mild aqueous conditions. Hybrid chemo-enzymatic routes may enable access to bioactive compounds not accessible by either approach alone.
5. **Circular economy for catalysts.** Recovery of catalysts from industrial waste streams and regeneration of deactivated catalysts are under-researched areas with significant environmental and economic potential.
6. **Green metrics for biological activity.** Current metrics focus on chemical efficiency. Incorporating biological activity (e.g., activity per unit waste) into sustainability assessment would better capture the therapeutic value of synthetic outputs.

References

1. Wiley Online Library. (2020). Advances in Unconventional and Sustainable Synthetic Approaches to Benzopyran Scaffolds. *The Chemical Record*, e70219.
2. Zenodo. (2020). Green Chemistry in Organic Synthesis: Principles, Methods, and Metrics for Sustainable Drug Development.
3. Europe PMC. (2020). A green fabrication of pharmacologically relevant fused pyrimidines using recyclable caffeine-H_3PO_4 catalyst: insight into antibacterial and cytotoxic efficacy.
4. ScienceDirect. (2020). Synthesis, characterization and catalytic activity of GO/ZnO nanocomposite for green synthesis of dihydropyrimidinones/thiones as potential antibacterial agents. *Inorganic Chemistry Communications*.
5. ScienceDirect. (2020). Benzethonium chloride:glycerol (1:2) deep eutectic solvent promoted green synthesis of benzo[4,5]imidazo[1,2-a]pyrimidine-3-carboxylate and evaluation of its anti-diabetic activity. *Synthetic Communications*.
6. Chemistry Europe. (2020). Ultrasound-Assisted Synthesis of 1,8-Dioxo-octahydroxanthenes Using N-Methylbenzimidazolium-Based Merrifield Resin-Supported Ionic Liquids as Recyclable Catalysts. *ChemistrySelect*.
7. RSC Publishing. (2020). Sustainable route to antiviral furano-chalcones via microwave-assisted solvent-free synthesis with recyclable MgO. *Green Chemistry*, 28, 5334.
8. RSC Publishing. (2020). Green and sustainable nanocatalyzed syntheses of hybrid heterocyclic scaffolds with privileged substructures. *Organic & Biomolecular Chemistry*, 24, 2189–2218.
9. ScienceDirect. (2020). Mechanistic and docking analysis of caffeine–H_3PO_4 catalyzed fused pyrimidine synthesis.
10. OUCI. (2020). An immobilized Schiff base–Mn complex as a hybrid magnetic nanocatalyst for green synthesis of biologically active [4,3-d]pyrido[1,2-a]pyrimidin-6-ones. *Nanoscale Advances*, 10, 2713–2721.
11. RSC Publishing. (2020). A Brønsted-based deep eutectic solvent as a green catalyst for the sustainable one-pot synthesis of 1,2,4,5-tetrasubstituted imidazoles. *RSC Advances*.

Research Paper © 2012 IJFANS. All Rights Reserved

12. ACS Publications. (2019). Recent Advances in Pyrimidine Molecular Hybrids, Green Synthetic Strategies, and Diverse Biological Applications. *ACS Omega*, 11(28), 41054.
13. Chemistry Europe. (2020). Catalyst Free, Microwave-assisted Synthesis of Benzylidene Schiff Bases and Evaluation of Their Biological Properties. *ChemistrySelect*, 9(43), e202404710.
14. Current Green Chemistry. (2020). A Review on Green Synthesis of Biologically Active Compounds. 9(3), 174–195.
15. Anastas, P. T., & Warner, J. C. (1998). *Green Chemistry: Theory and Practice*. Oxford University Press.
16. Clark, J. H., & Macquarrie, D. J. (Eds.). (2002). *Handbook of Green Chemistry and Technology*. Blackwell Science.
17. Dunn, P. J., Wells, A. S., & Williams, M. T. (Eds.). (2010). *Green Chemistry in the Pharmaceutical Industry*. Wiley-VCH.
18. Sheldon, R. A. (2017). The E factor 25 years on: The rise of green chemistry and sustainability. *Green Chemistry*, 19(1), 18–43.
19. Sheldon, R. A., & Woodley, J. M. (2018). Role of biocatalysis in sustainable chemistry. *Chemical Reviews*, 118(2), 801–838.
20. Trost, B. M. (1991). The atom economy—A search for synthetic efficiency. *Science*, 254(5037), 1471–1477.
21. Zhang, Q., De Oliveira Vigier, K., Royer, S., & Jérôme, F. (2012). Deep eutectic solvents: Syntheses, properties and applications. *Chemical Society Reviews*, 41(21), 7108–7146.