

GC-MS Analysis of the Phytochemical Constituents, and Anti-Inflammatory Activities, Anti-bacterial and Anti oxidant of *Nyctanthes Leaf Arbor-Tristis* Linn.

Ponnusamy Kiruba*¹, Kuppusamy Palanisamy², Ponnuswamy Selvamaleeswaran³,

¹Department of Botany, Arignar Anna Government Arts and Science College, Namakkal, India

²Department of Botany, Arignar Anna Government Arts and Science College, Namakkal, India

³Department of Biotechnology, Muthayammal College of Arts and Science, Rasipuram, India

Corresponding author: P. KIRUBA

corresponding author Mail ID: pkirubabotany006@gmail.com

Abstract:

Nyctanthes arbor-tristis is a member of the Oleaceae family with various therapeutic characteristics. Different plant components are used in traditional medicine to treat illnesses such as sciatica, chronic fever, and skin diseases. It is today regarded as a rich source of numerous unique compounds for creating medicines against various diseases and certain industrial products. In general, the leaves, roots, flowers, and seeds of Parijat are used to cure several ailments in various dose forms, including juice and powder. The current paper discusses the Plant Profile of *N. arbor-tristis* Linn., as well as the Phytochemical Screening of Extract of *N. arbor-tristis* Leaves, physicochemical parameters, and the Extraction Procedure and Pharmacological Activities conducted on *N. arbor-tristis* Leaves. The study aims to analyze the antioxidant, antibacterial, and anti-inflammatory potential of petroleum ether extract of *N. arbor-tristis* leaves. DPPH scavenging assays and *in vitro* anti-inflammatory activity was evaluated using Albumin denaturation, Proteinase inhibitory assay and was subjected to GC-MS analysis to identify its active components. The antibacterial potential was analyzed using the agar well diffusion method against Gram-negative and Gram-positive bacteria. Preliminary phytochemical screening confirmed the presence of alkaloids, flavonoids, phenols, tannins, triterpenoids, Amino acids, Carbohydrate, and the absence of alkaloids, Anthraquinone, Phenols, Saponin, and tannin. Remarkably, free radicals scavenging ability was observed in DPPH with the IC₅₀ of 610.71±13.6µg/mL, and the anti-inflammatory assay also observed Albumin denaturation with the IC₅₀ of 781.93±13.6µg/mL and the proteinase inhibitory activity with the IC₅₀ of 652.98±32.5µg/mL. In the antibacterial assay, the zone of inhibition was recorded as 20.6±1.6mm, 21.3±1.8mm, 21.3±1.8mm, and 17.9±1.5mm for the strains of *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas* spp., *Staphylococcus aureus*, and *Bacillus* spp. respectively. The highest inhibition zone was observed against *E. coli* (24.8±1.7 and 26.6±1.4mg/mL, respectively), and the least inhibitory was observed against *L. acidophilus* (19.5±1.1 and 21.7±1.7mg/mL) respectively. GC-MS profile exhibited the presence of 29 components highly accountable for its pharmaceutical activities. The results suggested the presence of phytochemical in

Nyctanthes arbor-tristis Leaves possess antioxidant and anti-inflammatory properties in the petroleum ether extract. This information can be used as a starting point in the search for natural-based drugs that are effective at alleviating inflammatory symptoms and antioxidant aspects.

Keywords: *Nyctanthes arbor-tristis* L, Antioxidant, Antibacterial, Anti-inflammatory, DPPH, Phytochemical profile, and GC-MS Analysis.

INTRODUCTION

Plants are natural synthesizers that are essential for meeting humanity's daily needs. Out of the 4, 22, 000 flowering plants reported worldwide, more than 50,000 are reported to have medicinal and pharmacological use. India contains a diverse range of medicinal plants¹⁻². Plants have long been used as medicines, according to ancient literature. Many major medications of the modern era have been made possible by such documentation of the most important traditional information on medicinal plants. The Charaka Samhita (1000 B.C.), our earliest text on Indian medicine, has a collection of information regarding the benefits of herbal medicines and references the utilization of over 2000 distinct plants³. According to a WHO report, 80% of the population in developing countries rely almost totally on traditional medicine for their primary healthcare needs⁴. Based on this research into active plant chemicals and pharmacological screening, researchers may be able to discover leads for the production of novel medications. So yet, just 6% of the estimated 400,000 plant species have had their biological activity investigated⁵. *N. arbor-tristis*, commonly known as Harshingar, coral Jasmine, Night Jasmine, and the queen of the night. The genus name 'Nyctanthes' is derived from two Greek words Nykhta (night) and Anthos (flower), and is consequently known as the night flower⁶ It is also familiar as "The sad tree" or "The tree of sorrow" since the blossoms bloom only during the night and are shed throughout the day. *N. arbor-tristis* leaf has shown promising results as psourcel source for its bioactive compounds with multipotent active prevents and reduces risk factors for various dis enhances enhancing certain physiological functions⁷⁻⁸. The disease-preventing nature is due to the presence of antioxidants, such as flavonoids, hydrolyzable tannins, phenolic acids, etc. These react against free radicals caused by various biological and environmental sources caused by an imbalance of naturally occurring antioxidants. It further contributes to developing diseases⁹ responsible for cancer, coronary heart diseases, and mutagenic and associated inflammation reactions¹⁰. Inflammation is frequently associated with pain and involves several occurrences, such as increased vascular permeability, protein denaturation, and membrane alteration. These occurrences can result in the loss of protein secondary and tertiary structure due to external stress or exposure to compounds like strong acids, bases, concentrated inorganic salts, organic solvents, or heat [11]. The immune modulation effect of anti-inflammatory agents includes modulating cytokine production and pro-inflammatory gene expression¹². They effectively treat infectious diseases and alleviate many side effects often associated with synthetic drugs¹³. Numerous computational methods are used to discover novel and highly effective drug candidates and promising targets in various diseases¹⁴.

Knowledge of the chemical constituents of plants is desirable not only for the discovery of therapeutic agents but also because such information may be of great value in disclosing new sources of economic phytochemicals for synthesizing complex chemical substances and discovering the actual significance of folkloric remedies¹⁵. Only very few reports have been published on *N. arbor-tristis* leaf extracts, and no systematic work has been done on the phytochemical and antioxidant potential; hence the present study has been taken with the aim of extraction, phytochemical characterization, and evaluation of antimicrobial and antioxidant potential.

MATERIALS AND METHODS

Collection of plant material

The leaves of *Nyctanthes arbor-tristis* were collected from Kolli Hills, Namakkal district, South India (Lat:11.2485°N; Long:78.3387°E). Dr. P. Subramaniam, Plant Taxonomist, Research Center in Botany, Aringar Anna Government Arts College, Namakkal, authenticated the plant. The voucher specimens (AGAR/BOT/001/2022 - 2023) were deposited in the Department of Botany, Aringar Anna Government Arts College, Namakkal, Tamil Nadu, India. Local Tamil names of the plants - Pavizhamalli or Parijata. The leaves were washed thoroughly with running tap water and distilled water, then shade-dried for 2-3 days, then oven dried and ground into a coarse powder using pestle-mortar. To prepare the petroleum ether extract, 30 g of a dried leaf was extracted in 300 mL of absolute petroleum ether using the Soxhlet apparatus for 8 hrs. The extracts were concentrated, stored at 4°C in a rotary evaporator, and freeze-dried for further use.

Preliminary phytochemical studies

The petroleum ether extract of *N. arbor-tristis* leaves was tested against microorganisms such as *Lactobacillus acidophilus*, *Escherichia coli*, *Salmonella typhi*, and *Klebsiella pneumonia*. The bacterial strains were maintained on nutrient agar slants at 4°C. A loopful of each bacterial strain was inoculated into 50mL sterile nutrient broth. The flasks were incubated at 37°C for 24 hrs.

Phytochemical screening

Preliminary phytochemical analysis was carried out for the extracts as per standard methods¹⁶

Detection of Total Alkaloids Content

Mayer's test: Filtrates were treated with Mayer's reagent. The formation of a yellow cream precipitate indicates the presence of alkaloids.

Detection of Flavonoids

H₂SO₄ test: Extracts were treated with a few drops of H₂SO₄. The formation of orange color indicates the presence of flavonoids.

Detection of Steroids

Liebermann- Burchard test: 2mL of acetic anhydride was added to 0.5g of the extracts, each with 2mL of H₂SO₄. The color changed from violet to blue or green in some samples, indicating the presence of steroids.

Detection of Terpenoids

Salkowski's test: 0.2g of the extract of the whole plant sample was mixed with 2mL of chloroform, and concentrated H₂SO₄ (3mL) was carefully added to form a layer. A reddish-brown coloration of the

inner face indicates the presence of terpenoids.

Detection of Anthroquinones

Borntrager's test: One gram (1 g) of the powdered seed was placed in a dry test tube, and 20 mL of chloroform was added. It was heated in a steam bath for 5 min. The extract was filtered while hot and allowed to cool. The filtrate was added with an equal volume of 10% ammonia solution. It was shaken, and the upper aqueous layer was observed for bright pink coloration to indicate the presence of Anthraquinones. In a test tube, control tests were done by adding 10 mL of 10 % ammonia solution in 5 ml chloroform.

Detection of Phenols

Ferric chloride test: Extracts were treated with a few drops of 5% ferric chloride solution. The formation of bluish-black color indicates the presence of phenol.

Detection of Saponins

Froth test: About 0.2g of the extract was shaken with 5 mL of distilled water. Formation of frothing (appearance of creamy stable persistent of small bubbles) shows the presence of saponins.

Detection of Tannins

Ferric chloride test: A small extract was mixed with water and heated in a water bath. The mixture was filtered, and 0.1% ferric chloride was added to the filtrate. A dark green color formation indicates the presence of tannins.

Detection of Carbohydrates

Fehling's test: 0.2gm filtrate is boiled in a water bath with 0.2mL each of Fehling solutions A and B. A red precipitate indicates the presence of sugar.

Detection of Oils and Resins

Spot test: Test solution of as little as 1 ng in a drop (ca. 50 μ L) of the solution was applied on filter paper. It develops a transparent appearance on the filter paper. It indicates the presence of oils and resins.

Gas Chromatography-Mass Spectrometry (GC– MS) Analysis

Gas Chromatograph-Mass Spectrometer quantitatively performed the GC-MS analysis of petroleum ether leaf extract. Hp5-MS column was used. High pure helium was the carrier gas, and the flow rate was 1ml/min. The temperature at the front inlet was 220°C. The oven temperature was maintained at 50°C to 250°C, gradually raised at 10°C/min. The ion chamber and GC interface temperature were maintained at 250°C. The identification of the peaks was based on computer matching of the mass spectra with the National Institute of Standards and Technology (NIST database) library and by direct comparison with published data¹⁷.

In vitro Antioxidant Activities

DPPH Free Radical Scavenging Activity

The plant extract at different concentrations was diluted with DMSO to get a sample solution. 5 μ L of the sample was seeded in a 96-well plate, followed by 195 μ L DPPH working solution to each well. After 20 min reaction, the absorbance was measured at 515 nm. The free radical activity of the extracts was determined by comparing their absorbance with blank¹⁸. The scavenging ability of DPPH radical was expressed as percentage inhibition and

calculated ($\mu\text{g/mL}$).

DPPH scavenging activity (%) = $(A_0 - A_1) / A_0 \times 100$

Where A_0 - absorbance of the control, and A_1 - absorbance of the sample.

In vitro anti-inflammatory activity of N. arbor-tristis

Inhibition of albumin denaturation

The anti-inflammatory activity of the sample extract was studied by using the inhibition of albumin denaturation technique which was followed with minor modifications¹⁹⁻²⁰. The reaction mixture consisted of test extracts and a 1% aqueous solution of bovine albumin fraction. pH of the reaction mixture was adjusted using a small amount of 1NHCl. The sample extracts were incubated at 37°C for 20 minutes and then heated to 51°C for 20 minutes. After cooling the samples, the turbidity was measured at 660 nm. (UV Visible Spectrophotometer Model 371, Elico India Ltd) The experiment was performed in triplicate. The Percentage Inhibition of protein denaturation was calculated as follows:

Percent inhibition (%) = $(\text{OD of Sample} - \text{OD of Control}) / \text{OD of Control}$
X 100.

Inhibition of Antiproteinase action

The reaction mixtures (2 mL) contained 0.06 mg trypsin, 1 mL 20 mM Tris HCl buffer (pH 7.4), and 1 mL test samples, each of different concentrations (100 - 500 $\mu\text{g/mL}$). The mixtures were incubated at 37°C for 5 min, and then 1 mL of 0.8% (w/v) casein was added. The mixtures were incubated for an additional 20 min. 2 mL of 70% perchloric acid was added to arrest the reaction. The cloudy suspension was centrifuged, and the absorbance of the supernatant was read at 210 nm against the buffer as blank. The experiment was performed in triplicate. The percentage inhibition of proteinase inhibitory activity was calculated as follows.

Percentage inhibition = $(\text{Abs control} - \text{Abs sample}) \times 100 / \text{Abs control}$

RESULTS AND DISCUSSION

Phytochemical Screening


Because of their many pharmacological qualities, plants are considered natural resources. Indeed, many phytochemicals, such as alkaloids, terpenoids, phenols, tannins, and so on, can be related to pharmacological activities. The primary metabolites include carbohydrates, fats, proteins, and minerals; phytoconstituents such as phenols and flavonoids are secondary metabolites generated from the primary metabolites. Studies in the literature have linked the existence of secondary metabolites to the biological capacity of plants²¹. Phytochemical screening of *N. arbor-tristis* leaves petroleum ether extract confirmed the presence of flavonoids, steroids, terpenoids, and carbohydrates oils & resins in different qualitative ranges. The negative sign depicted the absence of alkaloids, Arthroquinone, Phenols, Saponin, and tannin. The nine phytoconstituents, steroids, and terpenoids showed a strong presence depicting the pharmaceutical property of the *N. arbor-tristis* leaves aqueous extract. Hence all these secondary metabolites in the extracts were quantified further to assay the best extraction possessing the significant levels of bioactive components (Table 1 & Figure 1). In *N. arbor-tristis*, screening

of phytochemicals in methanol and aqueous extracts revealed phytochemicals such as phenols, terpenoids, and tannins only in the methanolic extract. In contrast, alkaloids and steroids were only present in the aqueous extract obtained through the cold percolation technique²². Preliminary phytochemicals such as carbohydrates, alkaloids, cardiac glycosides, phenol, flavonoids, and physicochemical parameters were also encountered²³. GC – MS results of petroleum ether extract leaves of *N. arbor-tristis* confirmed the presence of 29 components highly responsible for its antioxidant, antibacterial, antioxidant, and anti-cancer properties. The active biomolecules and their retention time (RT), peak area, molecular formula, molecular weight (MW), and structures obtained from PubChem sources are presented in Table 2. The first compound identified with less RT (4.342 min) was assigned as o-Xylene and the compound which took long RT (22.451min) was identified as Heptadecane, 2, 6,10, 15-tetramethyl. Specifically, Vitamins, Sterols, and their derivative compounds possess a high percentage of peak area among the entire phytochemicals (Figure 2 & Table 2). The compounds elucidated in GC–MS spectra are known for their therapeutic properties³⁴. GC-MS analysis of the methanolic extract from *N. arbor-tristis* L. leaves revealed the active compounds: Astragaloside, Nicotiflorin, Nyctanthic acid, Friedelin, and Lupeol³⁵. The study also supported the data, including the phytoconstituents such as D- mannitol, essential oil, and methyl salicylate. At the same time, the seeds contain glycerides of stearic, palmitic, oleic, linoleic, lignoceric, and myristic acids, lauric acid, D-glucose, and D- mannose. According to their reports, flower oil also includes 1-deconal anisaldehyde, Frie-del-1-ene-3-one, 1hydroxyfriedel-4-(23)-en-3-one, 1-triacontanol, lignoceric acid, and pelargonic acid were extracted from *N. arbor-tristis* stems. Sigmasterol and n-Tetradecyl-b-D-glucopyranoside, a newly found rangy-olone. There are also additional compounds, such as 2-phenyl ethyl b-D-glucopyranoside. Some natural compounds collected from this plant species have been reported to improve various biological conditions. Some of these discovered and extracted compounds have been demonstrated to heal many biological diseases³⁶.

Table: 1 Phytochemical Analysis of *N. arbor-tristis*

Table: 1 Phytochemical Analysis of <i>N. arbor-tristis</i>		
Phytochemicals	Observations	Inference
Alkaloids		-
Mayer's test	Cream color	
Wagner's test	Reddish brown solution/ precipitate	
Flavonoids		-
Lead acetate test	Yellow-orange	
H ₂ SO ₄ test	Reddish brown / Orange color precipitate	
Steroids	Violet to blue or green color formation	+
Liebermann-Burchard test		
Terpenoids		+
Salkowski test	Reddish brown precipitate	
Arthroquinone		-
Borntrager's test	Pink color	

Phenols		-
Ferric chloride test	Deep blue to Black color	
Lead acetate test	formation White precipitate	
Saponin	Stable persistent	-
Tannin	Brownish green / Blue black	-
Carbohydrates	Yellow/brownish/blue/green color	+
Oils & Resins	Filter paper method	+

Fig: 1 phytochemical analysis of *N. arbor-tristis*

In vitro antioxidant Activity

DPPH assay

GC-MS Analysis

Table: 2 Phytocomponents of *Nyctanthes arbor-tristis* Leaves extract by GC-MS

R/T	Name of the Compound	Molecular Formula	M/W	Molecular Structure	Peak Area%	Biological Activity
4.342	o-Xylene	C ₈ H ₁₀	106.16		1.12 %	Anti-proliferative

5.398	Cyclotetrasiloxane, octamethyl-	$C_8H_{24}O_6$ Si ₅	356.7	1.35 %	Antimicrobial & antioxidant
6.198	Ethyl 4-(5-methyl-1,1- dioxide- 4-	$C_{19}H_{19}$ NO ₅ S	373.4	5.68 %	Analgesic
6.842	1-(2,4- Dimethoxyphenyl)- propan-2	$C_{11}H_{14}$ O ₃	194.23	21.51 %	Antimicrobial
7.087	Undecane	$C_{11}H_{24}$	156.31	1.62 %	Cytotoxic
8.375	Diethylene glycol mono- dodecyl ether	$C_{16}H_{34}$ O	274.44	3.51 %	Anti-bacterial
9.009	Phenazine, 2-nitro-	$C_{12}H_7N$ 3O ₃	241.2	5.41 %	Anti-bacterial
10.28 6	Propenone, 3-(2- benzoxazolylthio	$C_{16}H_{11}$ NO ₂ S	281.3	2.19 %	Antimicrobial & Anticancer
10.731	7H- Dibenzoc,h]phenothiazine	$C_{20}H_{13}$ NS	299.4	2.42 %	Anticancer
10.875	Silanol, trimethyl-, phosphate	$C_9H_{27}O_4$ PSi ₃	314.54	2.91 %	Antioxidant & Antimicrobial
12.11 9	Octadecane, 1-bromo-	$C_{39}H_{79}$ BrO ₂	659.9	1.11 %	Antitumor
12.586	2-Methoxybenzoylformic acid, TM S	$C_{12}H_{16}O$ 4Si	252.34	4.37 %	Anticancer

Table: 2 Phytocomponents of *Nyctanthes arbor-tristis* Leaves extract by GC-MS

R/T	Name of the Compound	Molecular Formula	M/W	Molecular Structure	Peak Area%	Biological Activity
4.342	o-Xylene	C ₈ H ₁₀	106.16		1.12 %	Anti-proliferative
5.398	Cyclotetrasiloxane, octamethyl-	C ₈ H ₂₄ O ₆ Si ₅	356.7		1.35 %	Antimicrobial & antioxidant
6.198	Ethyl 4-(5-methyl-1,1-dioxide-4-	C ₁₉ H ₁₉ NO ₅ S	373.4		5.68 %	Analgesic
6.842	1-(2,4-Dimethoxyphenyl)-propan-2	C ₁₁ H ₁₄ O ₃	194.23		21.51 %	Antimicrobial
7.087	Undecane	C ₁₁ H ₂₄	156.31		1.62 %	Cytotoxic
8.375	Diethylene glycol mono-dodecyl ether	C ₁₆ H ₃₄ O	274.44		3.51 %	Anti-bacterial
9.009	Phenazine, 2-nitro-	C ₁₂ H ₇ N 3O ₃	241.2		5.41 %	Anti-bacterial
10.28 6	Propenone, 3-(2-benzoxazolylthio	C ₁₆ H ₁₁ NO ₂ S	281.3		2.19 %	Antimicrobial & Anticancer
10.731	7H-Dibenzoc,h]phenothiazine	C ₂₀ H ₁₃ NS	299.4		2.42 %	Anticancer
10.875	Silanol, trimethyl-, phosphate	C ₉ H ₂₇ O ₄ PSi ₃	314.54		2.91 %	Antioxidant & Antimicrobial
12.11 9	Octadecane, 1-bromo-	C ₃₉ H ₇₉ BrO ₂	659.9		1.11 %	Antitumor

12.586	2-Methoxybenzoylformic acid,	C ₁₂ H ₁₆ O ₄ Si	252.34	4.37 %	Anticancer
	TM				
	S				




Fig: 2 GC–MS chromatogram of *N. arbor-tristis*

Plants contain powerful antioxidants, and the application of *N. arbor-tristis* leaves as a decoction in Ayurveda medicine for a variety of diseases has been connected to the anti-oxidant activity of groups of chemicals that help in scavenging the free radicals that are primarily responsible for pathogenesis. Various experimental setups such as DPPH ((1,1-diphenyl-2-picrylhydrazyl), hydroxyl and superoxide radicals, and H₂O₂ scavenging studies demonstrate the exceptional antioxidant activity of *N. arbor-tristis*. The free radical scavenging activity of different concentrations of *N. arbor-tristis* leaves extracts and of standard ascorbic acid was evaluated by using DPPH radical scavenging method²⁷ *in vitro* DPPH assay for the plant extract *N. arbor-tristis* were analyzed by different concentrations (50, 250, 500, 750 and 1000 μ L). Inhibitory Concentration (IC₅₀) values were calculated. The inhibition concentration for the different plant concentrations was found as 31.72 \pm 2.7, 40.00 \pm 3.8, 48.97 \pm 3.4, 55.17 \pm 4.6 and 59.31 \pm 4.2% for the 50, 250, 500, 750, and 1000 μ L tested groups respectively. Finally, the overall inhibitory concentration of the plant *N. arbor-tristis* for DPPH assay is analyzed as 610.71 \pm 13.6 μ g/mL, increased concentration of plant extract showed increased scavenging activities against the reactive active species (Table 3 & Figure 2). The DPPH test of the methanolic extract revealed an inhibition percent of 69.42 at a 100g/mL concentration. However, aqueous extract only showed a 31.14 percent inhibition at the same concentration. Methanolic extracts were mostly effective at scavenging free radicals. The antioxidant potential data demonstrated inhibition in a dose-dependent way. The DPPH experiment revealed that the methanolic extract inhibited free radicals the most, whereas the aqueous extract inhibited them the least²¹. Various studies of free radical scavenging activity of various solvent extracts of dry and fresh flowers using multiple techniques, including such lipid peroxidation test, reducing activity, and H₂O₂ scavenging Compared to the control, all the treated groups showed high antimicrobial activity. In the control group, the highest microbicidal activity is exhibited against *E. coli* at 20.6 \pm 1.6mm, whereas the least micro-bacterial activity was

observed against *S. typhi* at 16.4 ± 1.7 mm (Figures 4 & 5). In the 40 mg/mL group, high inhibition was found against *E. coli* at 21.3 ± 1.8 mm, and the least inhibition was found against *S. typhi* at 17.9 ± 1.5 mm. In 60 assay, as well as changing amounts of enzymatic and non-enzymatic antioxidants, discovered that methanol extracts of dry flowers possess total phenolics content and antioxidant activity. In contrast, the aqueous extract of dry flowers has high enzymatic activity. Ethanol extracts of *N. arbor-tristis* stem exhibit concentration-dependent antioxidant activity in numerous animals. These data suggest that different antioxidants occur in the leaves, stems, and flowers of *N. arbor-tristis* 28.

S. No	Concentration (μ L)	IC ₅₀ (%)	IC ₅₀ (μ g /mL)
1	50	31.72±2.7	610.71±13.6
2	250	40.00±3.8	
3	500	48.97±3.4	
4	750	55.17±4.6	
5	1000	59.31±4.2	




Fig: 3 In vitro DPPH assay of *N. arbor-tristis****Antibacterial Activity of *N. arbor-tristis* leavespetroleum ether extract.***

Antimicrobial activities of the plant extract are the essential parameters for the active compound analysis in phytosources²⁹. By disc diffusion method, the antimicrobial activity of the plant extract *N. arbor-tristis* is analyzed against four bacterial (*L. acidophilus*, *E. coli*, *S. typhi*, and *K. pneumoniae*) species tested. Three concentrations (40, 60, and 80mg/mL) and control groups are maintained. and 80mg/mL groups, the highest inhibition zone are observed against *E. coli* (24.8±1.7 and 26.6±1.4mg/mL, respectively), and the least inhibition activity was observed against *L. acidophilus* (19.5±1.1 and 21.7±1.7mg/mL) respectively. Significantly increased concentration of plant extract showed significant inhibition against the testing microbes (Table 4).

Organisms	Control	Inhibition zone (in mm)		
		40mg/m L	60mg/m L	80mg/m L
<i>Lactobacillus acidophilus</i>	17.8±1.1	18.8±1.9	19.5±1.1	21.7±1.7
<i>Escherichia coli</i>	20.6±1.6	21.3±1.8	24.8±1.7	26.6±1.4
<i>Salmonella typhi</i>	16.4±1.7	17.9±1.5	19.8±1.5	21.9±1.6
<i>Klebsiella pneumoniae</i>	19.2±1.5	20.0±1.7	22.7±1.8	24.8±1.3

Anti-inflammatory Activity***Inhibition of albumin denaturation***

Denaturation of proteins is a well-certified cause of inflammation. As part of the investigation on the mechanism of the anti-inflammation activity, the capability of extract to inhibit protein denaturation was studied^{30 - 31}. Albumin denaturation inhibition activity for the plant extract *N. arbor-tristis* was analyzed by different concentrations (50, 250, 500, 750, and 1000µL). The inhibition concentration for the different plant concentrations was found as 26.97±2.85, 32.89±3.13, 41.45±3.51, 49.34±4.21 and 56.58±3.86% for the 50, 250, 500, 750, and 1000µL tested groups respectively. Finally, the overall albumin denaturation inhibitory concentration of the plant *N. arbor-tristis* is 781.93±13.6µg/mL. Increased concentration of plant extract showed increased inhibition of albumin denaturation activities (Table 5 & Figure 6).




Fig: 4 In vitro Antibacterial Activity of *N. arbor-tristis*




Fig: 5 Minimum Inhibitory Concentration of *N. arbor-tristis* extracts

S. No	Concentration (μ L)	IC50%	IC50 (μ g /mL)
1	50	26.97 ± 2.8 5	781.93 ± 13.6

2	250	32.89±3.1 3
3	500	41.45±3.5 1
4	750	49.34±4.2 1
5	1000	56.58±3.8 6




Fig: 6 *In vitro* Inhibition of albumin denaturation of *N. arbor-tristis*

Proteinase inhibitory activity

Neutrophils are a rich resource of serine proteinase and are localized at lysosomes. It was previously reported that leukocyte proteinase plays an important role in the development of tissue damage during inflammatory reactions. A significant level of proteinase was provided by proteinase inhibitors³², and the Petroleum ether extract of *N. arbor-tristis* whole plant exhibited significant anti-proteinase inhibition activity for the plant extract of *N. arbor-tristis* are analyzed by different concentrations (50, 250, 500, 750, and 1000µL). The inhibition concentration for the is found as 32.26±1.99, 41.29±2.31, 45.81±2.75, 53.55±2.41 and 58.06±1.90% for the 50, 250, 500, 750, and 1000µL tested groups respectively. Finally, the overall albumin denaturation inhibitory concentration of the plant *N. arbor-tristis* is 652.98±32.5µg/mL. Increased concentration of plant extract showed increased inhibition of anti-proteinase activities. (Table 6 & Figure 7).

Table 6: <i>in vitro</i> Inhibition of Anti-proteinase action of			<i>Nyctanthes arbor-tristis</i>
S. No	Concentration (µ L)	IC50%	IC50 (µ g/mL)
1	50	32.26±1.9	652.98±32.5

		9
2	250	41.29±2.3
		1
3	500	45.81±2.7
		5
4	750	53.55±2.4
		1
5	1000	58.06±1.9
		0




Fig: 7 In vitro Inhibition of Anti-proteinase action of *Nyctanthes arbor-tristis*

CONCLUSION

According to the results of *in-vitro* and *in-vivo* investigations, *N. arbor-tristis* is a potentially enormous therapeutic in the field of herbal medicine, notably in the management of diseases including malaria, fevers, arthritis, and protozoan infections such as leishmaniasis. It has also been established that every part of this plant has therapeutic potential. Despite using crude extract, the only research that has undergone clinical trials is for treating malaria. Though the potentiality may widen, the studies faced limitations in exploring their research aspects. Furthermore, more detailed safety data on acute, subacute, cardiac, and immunotoxicity for crude and purified extracts is required. The present study highlights the petroleum ether extracts of the leaf of *N. arbor-tristis* possess multipotent activities. It is a potential antibacterial agent against pathogenic bacteria and may help interact with various plant diseases and human allergies. *N. arbor-tristis* has a better inhibitory effect on trypsin and a good inhibitory effect on protein denaturation and good antioxidants. Wherein GC-MS analysis revealed, 29 components in *N. arbor-tristis* leaves petroleum ether extract. The plant material could be exploited as an alternative antibiotic drug discovery toward combating the global disease burden. The potential economic and social impact of the use of *N. arbor-tristis* in the management of various diseases. Thus, the importance of continued research in this area and the potential benefits that could be derived from herbal medicine.

ACKNOWLEDGEMENTS

The authors thank the tribal people near the Namakkal area for sharing their plant knowledge.

AUTHORS CONTRIBUTION STATEMENT

Ms. Ponnusamy Kiruba & Dr. Michael Helan Soundra Rani collected the samples from the Kolli Hills, Namakkal district, and sketched the concepts in processing & extraction procedures. Dr. Ponnuswamy Selvamaleeswaran designed and performed the anti-inflammatory activity and GCMS characterization study. Dr. Edward Gnanaraj Wesely and Dr. Kuppusamy Palanisamy analyzed the data regarding Methodology & Validate the original draft writing with suggestions and inputs from all authors.

REFERENCES

1. Veeresham C. Natural products derived from plants as a source of drugs. *J Adv Pharm Technol Res.* 2012; 3(4):200-1. doi: 10.4103/2231-4040.104709, PMID 23378939.
2. Dias DA, Urban S, Roessner U. A historical overview of natural products in drug discovery. *Metabolites.* 2012; 2(2):303-36. doi: 10.3390/metabo2020303, PMID 24957513.
3. Shi Y, Zhang C, Li X. Traditional medicine in India. *J Trad Chin Med Sci.* 2021;8:S51-5. doi: 10.1016/j.jtcms.2020.06.007.
4. Tamta A, Tailor CS. Pharmacognostic, ethnomedicinal, phytopharmacological review of *Oxalis corniculata* Linn.
5. shetty P. S, Venkatesh MD. Scope and implementation strategies to streamline medical pluralism in the contemporary healthcare system. *J Res Ayurvedic Sci.* 2023;7(1):30.
6. Bhadra P. Efficacy of some compounds isolated from *Nyctanthes arbor-tristis* Linn. on human and plant diseases revealed in silico analysis. *Indian J Nat Sci.* 2020;10(60):20833-9.
7. Singh J, Singh AP, Singh AP. *Nyctanthes arbor-tristis*: a comprehensive review. *World J Curr Med Pharm Res.* 2021;74-8. doi: 10.37022/wjcpr.v3i4.181.

8. Mugaranja KP, Kulal A. Alpha-glucosidase inhibition and antioxidant activity of *Ensete superbum* (Roxb.) Cheesman seeds: GC-MS-based profiling of the active metabolites and molecular docking study. *Curr Sci.* 2022;122(6):682-8. doi: 10.18520/cs/v122/i6/682-688.
9. Sigdel B, Prasad Khanal D, Raj Dhungana B, Lama D. Phytochemical and Biological study on leaves extract of *Nyctanthes arbor-tristis* L. *Asian J Pharmacogn.* 2021;5(3):5-14.
10. Ololade ZS, Kuyooro SE, Ogunmola OO, Abiona OO. Phytochemical, antioxidant, antiarthritic, anti-inflammatory, and bactericidal potentials of the leaf extract of *Lactuca teraxacifolia*. *Glob J Med Res B Pharm Drug Discov Toxicol Med.* 2017;17:19-27.
11. Talukdar R, Wary S, Mili C, Roy S, Tayung K. Antimicrobial secondary metabolites obtained from endophytic fungi inhabiting healthy leaf tissues of *Houttuynia cordata* Thunb., an ethnomedicinal plant of Northeast India. *J Appl Pharm Sci.* 2020;10(09):099- 106.
12. Amadarshanie DBT, Gunathilaka TL, Silva MR, Navaratne SB, Dinithi L, Peiris C. Functional and antiglycation properties of cow milk set yogurt enriched with *Nyctanthes arbor-tristis* L. flower extract. *Food Sci Technol.* 2022;154:1-9.
13. Tallei TE, Fatimawali YA, Marfuah S, Tania AD, Kalalo MJ, Johnson Kepel B, Nurdjannah Jane Niode N, Kusumawaty D, Idroes R. Evaluation of the Potential for Immunomodulatory and anti-inflammatory Properties of phytoconstituents Derived from Pineapple *Ananas comosus* (L.) Merr. Peel Extract Using an in-silico Approach. *Philipp J Sci.* 2022;151:397- 410.
14. Jose D, Pandiammal S, Senthilkumaar P. Phytochemical Screening, antimicrobial and Antioxidant Potential of *Nyctanthes arbor-tristis* L. Floral extracts. *J Acad Ind Res.* 2016;5:35-9.
15. Sundararaj JP, Kuppuraj S, Ganesan A, Ponnusamy P, Nayaka S. in vitro assessment of antioxidant and antimicrobial activities of different solvent extracts from lichen *Ramalina nervulosa*. *Int J Pharm Pharm Sci.* 2015;7:200-4.
16. Brain KR, Tuner TD. The practical evaluation of phytopharmaceuticals. Bristol: Wright Scientectica Publishers; 1975. p. 57-8.
17. Sandhya Kumari TD, Singara Charya MA. Phytochemistry, anticancer and anti-inflammatory activities of solvent leaf extracts of *Nyctanthes arbor-tristis*. *Int J Pharm Sci Res.* 2017;8(4):1654-63.
18. Molyneux P. The use of the stable free radical diphenyl picrylhydrazyl (DPPH) for

estimating antioxidant activity. Songlanakar J Sci Technol. 2004;26:211-9.

19.Rajesh A, Doss A, Tresina PS, Mohan VR. In vitro anti- inflammatory activity of ethanol extract of *Caralluma umbellata* haw (Asclepiadaceae). Res J Life Sci Bioinformatics Pharm Chem Sci. 2019;5(2):515.

20.Mizushima Y, Kobayashi M. Interaction of anti- inflammatory drugs with serum proteins, especially with some biologically active proteins. J Pharm Pharmacol. 1968;20(3):169-73. doi: 10.1111/j.2042- 7158.1968.tb09718.x, PMID 4385045.

21.Sakat SS, Juvekar AR, Gambhire MN. In vitro antioxidant and anti-inflammatory activity of methanol extract of *Oxalis corniculata* L. Int J Pharm Pharm Sci. 2010;2:146-55.

22.Kumar KD, Rani VP, Parimalam M, Kumar PPY, Venkadesh B. Preliminary pharmacological studies on *Nyctanthes arbor-tristis* L. flower extract. Ann Rom Soc Cell Biol. 2021;25(6):1759-69.

23.Daund AK, Jadhav RS, Vikhe DN. A Review on *Nyctanthes arbor-tristis* Leaves: A Potential Medicinal Herb, World. J Pharm Med Res. 2022;8(4):128-32.

24.Ekambaram H, Manjunath K, Udayashankara A. Antioxidant property and GC-MS profiling of the methanolic leaf extract of *Plumbago zeylanica*. Crit Rev. 2020;7:1428-37.

25.Pandrangi SL, Chalumuri SS, Chittineedi P, Garimella SV. Therapeutic potential of *Nyctanthes arbor-tristis* on cancer and various diseases. Ann Rom Soc Cell Biol. 1690-1701;26(01).

26.Michael JS, Kalirajan A, Padmalatha C, Singh AR. In vitro antioxidant evaluation and total phenolics of methanolic leaf extracts of *Nyctanthes arbor-tristis* L. Chin J Nat Med. 2022. 2013;11(5):484-7. doi: 10.1016/S1875-5364(13)60088-6, PMID 24359771.

27.Gulshan B, Suri KA, Parul G. A comprehensive review on *Nyctanthes arbor-tristis*. Int J Drug Deliv Res. 2015;7(1):183-93.

28.George S, Benny PJ, Kuriakose S, George C. Antibiotic activity of 2, 3-dihydroxybenzoic acid isolated from *Flacourtia inermis* fruit against multidrug-resistant bacteria. Asian J Pharm Clin Res. 2011;4:120-30.

29.Chandra Mishra R, Kumari R, Yadav S, Parkash Yadav J. GC-MS analysis and molecular docking studies of active phytochemicals from medicinal plants against *Malassezia globosa* LIP1 (SMG1) enzyme. Platinum open Access [journal]. 2022;12:2762-75.

- 30.Pandey AK, Kashyap PP, Kaur CD. Anti-inflammatory activity of novel Schiff bases by in vitro models. Bangladesh J Pharmacol. 2017;12(1):41-3. doi: 10.3329/bjp.v12i1.29675.
- 31.Udaya Prakash NK, Sriraman V, Ranjith Kumar M, Sripriya N, Bhuvanewari S. Antioxidant potency and GC-MS composition of leaves of Artocarpus altilis (Park). Fosh. Pharm Chem. 2017;9:102-6.
- 32.Prakash P, Bommana C, Nagaraj R, Shobana N, Uthralakshmi SS, Antony V. Samrot. Evaluation of Antioxidants, Antidiabetic, anti-inflammatory Active compounds from Leptogium rivurale through in vitro and in silico Studies. Platinum open Access [journal]. 2022;11:4192-200.