

CHECKLIST FOR FERN DIVERSITY OF WEST KARBI ANGLONG DISTRICT OF ASSAM, NORTHEAST INDIA

Utpal Phukan¹, Rakhal Roy Choudhury²

Department of Botany,

Rabindranath Tagore University, Hojai, Assam, 782435

Abstract: Pteridophytes are distributed in all the phytogeographical zones of India ranging from aquatic to alpine habitat where they grow as Hydrophyte, Mesophyte, Lithophyte, Epiphyte, Hemi epiphyte, Climbers etc. Present study finds the habitat of pteridophytes in riverine forests, forest floors, on slopes, grasslands, on rocks and crevices, on open walls and stone boulders etc. The study is carried on the area of West Karbi Anglong district of Assam in which major part of the area is hilly and some parts with plains and river valleys. The present study is focused on Fern diversity where 52 species were collected belonging to 31 genera under 20 families.

Keywords: Pteridophytic flora, Habitat, Species, Economic importance, Conservation, West Karbi Anglong.

INTRODUCTION

The first references to Indian Pteridophytes are in the classical Sanskrit linked to Ayurveda. In **Charak** and **Shushrut Samhitas Mayur Sikha** (Actiniopteris), Hansraj and Hanspadi (*Adiantum spp.*) were mentioned with medicinal properties. In Buddhist literature, it is thought that the species of Moonworts (*Botrychium spp.*) possess certain magical values and the plants of Botrychium are kept near the statue of Buddha to keep the devil powers away. Pteridophytes, also known as ferns and ferns-allies, flowerless plants without seeds and regarded as the first terrestrial plants to develop a vascular tissue system. Pteridophytes occupy an important position in the vegetation of the land and appeared for the first time around 360 million years ago marking the beginning of the Mesozoic era. They flourished on Earth for 200 million years before flowering plants developed. However, they do not form the dominant vegetation anywhere in the world today, but in few areas of rainforest they exhibit luxurious growth. They constitute an important group of plant diversity and are always present in moist shaded areas, often in abundance, but some species also reach xeric habitats. The main body of the plant is sporophyte, which is the dominant phase of its life cycle. Chapman (2009) estimated that there are about 12,000 species of ferns and fern-allies across the world. According to Moran (2015), “worldwide, there are about 13,600 species of ferns and lycophytes”. According to a survey, the pteridophytic flora of India comprises around 1,000 species belonging to 191 genera and 67 families including 47 endemic Indian ferns (Dixit, 1984) and in another report, more than 1,100 species of pteridophytes belonging to 144 genera and 34 families with about 235 endemic species (Chandra, 2000)

from India. According to Fraser-Jenkins *et. al.* (2017), “altogether there are about 1135 species including 42 exotics and 53 further subspecies”, from the Indian subcontinent. The north-eastern region of India, including the state Assam, is part of one of the world’s biodiversity hot spot regions, with a wide range of physiographic and climatic conditions. This area is being the home of many distinct ecological niches as the area is rich and diversified in flora. From the richness of its plant wealth and great diversity of vegetation, the northeast, formerly Assam in a wide sense, is often called the “**Floristic Gateway of India**”. The area is home to many early taxa and a wide range of endemic and threatened species. (Anonymous, 2006).

REVIEW OF LITERATURE

Previously, several workers had documented pteridophytes in north-eastern India. Among them, the oldest records are those of Kachroo (1953, 1975) who recorded ferns from various parts of the Assam. Panigrahi (1960), Panigrahi & Chowdhury (1961, 1962), Panigrahi & Patnaik (1961, 1968), etc. had listed the eastern India ferns from time to time. Over the past few years, many workers have conducted research into the species diversity of Pteridophytes in present time. Some of the important works are Dutta *et. al.* (1980) on the pteridophyte diversity of N. C. Hills district, Barua *et.al.* (1989) of Kamrup district, Bhattacharya (1994) of Karimganj district, Bhattacharya *et. al.* (1995) of Cachar district, Das *et. al.* (2012) in Nazira Subdivision of Sivasagar district etc. Bir *et. al.* (1992) published the North-Eastern Indian Pteridophyte species catalogue and Borthakur *et. al.* (2000) published a work on ferns and allied ferns in Assam. Chandra *et. al.* (2008) evaluated rare and threatened pteridophytes in India and found 414 species, 219 of which are at risk, 160 are critically endangered, 82 are near threatened and 113 are rare. However, this group of plants has not yet received enough attention due to the difficulty in identification and exploration in remote areas.



Figure : 1, Map showing the study area (West Karbi Anglong District of Assam, India. *Map not in scale*)

STUDY AREA

The Karbi Anglong district, the largest district of Assam, houses five wildlife sanctuaries viz., Nambor, East Karbi Anglong, North Karbi Anglong, Garampani, and Marat Longri Wildlife Sanctuary. East Karbi Anglong Wildlife Sanctuary (EKAWLS) is located in the eastern part of the Karbi Anglong district and is an integral part of the Karbi Anglong-Kaziranga landscape. The present study area is selected in the area of newly upgraded to district as West Karbi Anglong in 2016. Headquarter is Hamren. 70% of the area is hilly and covered by moist and mixed semi evergreen forest. Other part is composed of plain river valleys with human habitat and cultivated lands. The hilly terrain is continuous to the plateau of Meghalaya on west. The district is situated in the latitude of 25°33'N to 26°09'N and longitude of 92°08'E to 93°04'E. The climate is hot and humid, annual temperature ranges from 6 °C to 12°C and 23°C to 32°C in summer. The average rainfall is about 2416 mm. Annual precipitation of 1800 mm.

It is considered to be one of the rich floral and wildlife diversity regions of the Indo-Burmese biodiversity hotspot (WWF, 2002). The topography is varied with undulating hills, large valleys, rugged gorges, rivers and streams, annual and perennial streams and few waterfalls. The area is home to different types of forests ranging from semi-evergreen moist, moist mixed deciduous to various types with strewn patches of pure or mixed bamboo. The ground is well drained, sandy loam to clayey loam.

METHODS

A number of field visits were undertaken to the study area in different seasons to make a complete record of different habitats of the species and their occurrence during 2020-2021. Specimens were collected with their fertile and vegetative parts. Herbarium specimens were prepared by following standard herbarium technique (Jain & Rao 1977). Digital photographs of the concerned species were taken. Field notes viz. ecological data, habit, habitat and some diagnostic characters of each species were recorded. The specimens were kept in herbarium of department of Botany, RTU, further studied with standard literature like Baishya & Rao (1982), Jamir & Rao (1988), and Borthakur *et. al.* (2001, 2018). Botanical names were verified by consulting with the herbaria of Gauhati University and the Botanical Survey of India, Shillong (ASSAM). The nomenclature of the species has been given according to the International Plant Names Index (IPNI), available at www.ipni.org.

RESULTS & DISCUSSION

A total of 52 species of pteridophytes belonging to 31 genera and 20 families are recorded (Table 1). The Pteridaceae family had the highest number of species among the fern allies with 7 species (14.5%) belonging to 2 genera, which is followed by Thelypteridaceae family with 5 species (10.41%) from 1 genus. The third most dominant families are Lycopodiaceae, Polypodiaceae and Salviniaceae with 4 species each (8.3%). Besides these, the families Dryopteridaceae and Selaginellaceae has 3 species each (6.25%); Aspleniaceae, Athyriaceae, Blechnaceae, Davalliaceae, Dennstaedtiaceae, Lygodiaceae, Oleandraceae with 2 species (4.16%) each and Cyatheaceae, Equisetaceae, Gleicheniaceae and Marsileaceae with 1 species each (2.08%) each. This is the characteristic features of the vegetation of the region.

Table:1

Sl. No.	Family	Genus	Species	Habitat	Place of collection	Distribution
1.	Adiantaceae	<i>Adiantum</i>	<i>Adiantum incisum</i> Forssk.	1	Sekso Hill	Abundant
		<i>Adiantum</i>	<i>Adiantum philippense</i> L.	1	Umpanai	Frequent
		<i>Cheilanthes</i>	<i>Cheilanthes argentea</i> (Gmel.) Kunze.	1	Sekso Hill	Frequent
2.	Aspleniaceae	<i>Asplenium</i>	<i>A. nidus</i> L.	4	Dongka Sib Mandir	Abundant
		<i>Asplenium</i>	<i>A. phyllitidis</i> D. Don	4	Boithalangso, Zirikending	Abundant
3.	Athyriaceae	<i>Diplazium</i>	<i>D. dilatatum</i> Bl.	1	Sekso Hill, Zirikending	Abundant
		<i>Diplazium</i>	<i>D. esculentum</i> (Retz.) Sw.	1	Sekso Hill, Zirikending	Abundant
4.	Blechnaceae	<i>Blechnum</i>	<i>B. orientale</i> L.	1	Sekso Hill, Ronghang Rongbong	Abundant
		<i>Stenochlaena</i>	<i>S. palustris</i> (Burm.f.) Bedd.	5	Khanduli roadside	Abundant
5.	Cyatheaceae	<i>Cyathea</i>	<i>C. gigantea</i> (Wall. ex Hook.) Holttum	1	Near khanduliwater falls	Rare
6.	Davalliaceae	<i>Araiostegia</i>	<i>A. divaricata</i> (Bl.) M.Kato.	4	Khanduli	Abundant
		<i>Davallia</i>	<i>D. griffithiana</i> Hook. f.	4	Khanduli	Abundant
7.	Dennstaedtiaceae	<i>Microlepia</i>	<i>M. speluncae</i> (L.) T.Moore	1	Donka, Khanduli	Abundant
		<i>Odontosoria</i>	<i>O. chinensis</i> (L.) J. Sm	1	Amtring	Abundant
8.	Dryopteridaceae	<i>Bolbitis</i>	<i>B. heteroclita</i> (Presl.) Ching	1	Umpanai	Abundant
		<i>Tectaria</i>	<i>T. coadunata</i> (Wall. ex Hook. & Grev.) C. Chr	1	Umpanai	Abundant
		<i>Pleocnemia</i>	<i>P. submembranacea</i> (Hayata) Tagawa & K. Iwats.	1	Sekso Hill	Frequent
9.	Equisetaceae	<i>Equisetum</i>	<i>E. debile</i> D. Don	1	Donka Sib Mandir	Frequent
10.	Gleicheniaceae	<i>Dicranopteris</i>	<i>D. linearis</i> (Burm.f.) Underw.	1	On the way to Khanduli	Abundant
11.	Lycopodiaceae	<i>Huperzia</i>	<i>H. phlegmaria</i> (L.) Rothm.	4	Koka	Rare
		<i>Huperzia</i>	<i>H. squarrosa</i> (G.Forst.) Trevis	4	Umpanai	Rare
		<i>Lycopodiella</i>	<i>L. cernua</i> (L.) Pic. Serm	1	Koka	Frequent
		<i>Lycopodium</i>	<i>L. cernuum</i> L.	1	Sekso Hill	Frequent
12.	Lygodiaceae	<i>Lygodium</i>	<i>L. japonicum</i> (Thunb.)	5	Sekso Hill	Frequent
		<i>Lygodium</i>	<i>L. microphyllum</i> (Cav.) R. Br	5	Sekso Hill	Frequent

13.	Marattiaceae	<i>Angiopteris</i>	<i>Angiopterisevecta</i> (G. Frost.) Hoffm	1	Way to Kanduli from Hamren near waterfalls	Rare
14.	Marsileaceae	<i>Marsilea</i>	<i>M. quadrifolia</i> L.	2	Donka	Abundant
15.	Oleandraceae	<i>Nephrolepis</i>	<i>N. bisserata</i> (Sw.) Schott	4	Sekso Hill	Frequent
		<i>Nephrolepis</i>	<i>N. cordifolia</i> C. Presl.	1	Amtring	Frequent
16.	Pteridaceae	<i>Adiantum</i>	<i>A. capillus-veneris</i> L.	1	TikaPahar	Frequent
		<i>Adiantum</i>	<i>A. philippense</i> L.	1	Umpanai	Abundant
		<i>Pteris</i>	<i>P. ensiformis</i> Burm. f.	1	Khanduli water falls	Abundant
		<i>Pteris</i>	<i>P. longipinnula</i> Wall. ex J. Agardh.	1	Khanduli water falls	Abundant
		<i>Pteris</i>	<i>P. multifida</i> Roxb.	1	Amtring	Abundant
		<i>Pteris</i>	<i>P. semipinnata</i> L.	1	Amtring	Abundant
		<i>Pteris</i>	<i>P. biauritas</i> sub sp. <i>walkeriana</i> Fraser-Jenk. &Rajkumar	1	Umpanai	Abundant
17.	Polypodiaceae	<i>Microsorium</i>	<i>M. punctatum</i> (L) Copel.	4	Ulukunchi	Abundant
		<i>Pyrrosia</i>	<i>P. lanceolata</i> (L.) Farewell.	4	Tumpren, Donka	Abundant
		<i>Pyrrosia</i>	<i>P. nuda</i> (Giesenh.) Ching	4	Koka	Abundant
		<i>Pyrrosia</i>	<i>P. piloselloides</i> (L.) M.G.Price	4	Koka, Zirikending, Hamren	Abundant
18.	Salviniaceae	<i>Azolla</i>	<i>A. pinnata</i> R. Br. subsp. <i>asiatica</i> R.M.K. Saunders & K. Fowler	3	Tumpreng	Abundant
		<i>Salvinia</i>	<i>S. cucullata</i> Roxb.	3	Tumpreng	Abundant
		<i>Salvinia</i>	<i>S. molesta</i> D. S. Mitch.	3	Tumpreng	Abundant
		<i>Salvinia</i>	<i>S. natans</i> (L.) All.	3	Tumpreng	Abundant
19.	Selaginellaceae	<i>Selaginella</i>	<i>S. helferi</i> Warb.	1	Donka Sib Mandir	Frequent
		<i>Selaginella</i>	<i>S. monospora</i> Spring	1	Amtring	Frequent
		<i>Selaginella</i>	<i>S. semicordata</i> (Wall. ex Hook. Grev.) Spring	1	Boithalangso	Abundant
20.	Thelypteridaceae	<i>Thelypteris</i>	<i>T. dentata</i> (Forssk.) E.P.St.John	1	Sekso Hill	Abundant
		<i>Thelypteris</i>	<i>T. interrupta</i> (Willd.) K.Iwats.	1	Umpanai	Abundant
		<i>Thelypteris</i>	<i>T. ornata</i> (Wall ex Bedd.)	1	Sekso Hill	Abundant
		<i>Thelypteris</i>	<i>T. prolifera</i> (Retz.)	1	Hamren	Abundant

		C.F.Reed			
	<i>Thelypteris</i>	<i>T. triphylla</i> (Sw.) K. Iwats.	1	Sekso Hill	Abundant

1 – Terrestrial, 2 – Marshy Species, 3 – Free Floating, 4 – Epiphytic and 5 – Climbers

According to habitat-types the species were further classified under 5 broad groups viz., Terrestrial (33 species, 63.43%), Epiphytic (11 species, 21.15%) Free floating (4 species, 7.69%) Climbers (3 species, 5.76%) and Marshy places (1 species, 1.92%).Based on distribution, 35 species among total were Abundant (67.30%), 13 species were Frequent (25%) and 4 were rare (7.69%).




Plate: 1- Some photographs of collected specimens

1. *Adiantum incisum* Forssk. 2. *Adiantum philippense* L. 3. *Blechnum oriental* L.(Sporangium) 4. *Dicranopteris linearis* (Burm.f.) Underw. 5. *Cheilanthes argentea* (Gmel.) Kunze. 6. *Microlepia speluncae* (L) T. Moore

ECONOMIC IMPORTANCE

Humans have been depending on plants as an important source of medicine since ancient times. Vegetable curry is made with the tender leaves of *Dryopteris cochleata*, *D. sparsa* and *Leucostegia immersa*. *Diplazium esculentum* (Retz.) Sw.is used extensively as vegetables in the neighbouring region.The fresh leaves of *Diplazium esculentum* (Retz.) Sw., *Helminthostachy szeylanica* (L.) Hook., *Ophioglossum reticulatum* L. and *Marsilea minuta* L. are mixed with legumes and consumed with selection in north east India. Due to ornamental nature of fronds of *Dryopteris*, *Adiantum*, *Nephrolepis* etc., the potted plants of such species find place in terrace of houses, hotels and gardens etc. Dried stems are also used for ornamental purposes. The pith of *Angiopteris evecta*, *Cibotium assamicum* and tree ferns are being eaten by some tribals

in the period of shortage of food, as such or sometimes after colling also in the Eastern India and Arunachal Pradesh. *Adiantum philippense* L. and *Marsilea minuta* L. are referred to as important medications in 'Charak Samhita'. *Selaginell abryopteris* popularly referred to as **Sanjiwani** is widely used as a tonic in India. Lycopodium spores serve as a powder to the skin. The Lycopod spores known as vegetable brimstone are used for fireworks. The decoction of *Actiniopteris radiata* frond is used to control excessive bleeding in womens during mensuration or abortion. The decoction of *Adiantum lunulatum* used as diuretic, and useful in dysentery. The fronds of *Asplenium adiantum*, *A. nigrum* leaves are used in spleen disorder and also used among ladies for family planning in Unani system of Medicine. Rhizome of *Helminthostachys zeylanica* (L.) Hook. is used against dysentery, and leaves are used to cure sores on tongue. The stems of *Lygodium flexuosum* boiled with mustard oil and used against rheumatism. The rhizome of *Polystichum squamosum* known as 'Nirviri' in India, used against scorpion and insect bites. A juice is extracted from the rhizomes of *Tectaria macrodonta* in Darjeeling district and used against diarrhea among children. The leaf and root decoction of Frequently occurring *Adiantum philippensis* has been found to be very effective in the treatment of chest complaints. *Adiantum philippense* L. and *Lygodium microphyllum* (Cav.) R.Br., are used as a medicine in fever, dysentery, jaundice, etc. (Patiri & Borah, 2007; Sen & Ghosh, 2011). The fronds of *Pteris* Spp. are used to cure cuts and wounds. Orchids are grown in the main stems of *Osmunda*, *Alsophila*, *Angiosphila*, *Angiopteris*. There are many species of Pteridophytes which can be useful in the preparation of medicines. The medicinal qualities of ferns, real or imagined, are mentioned as far back as 300 B.C. by the Greek philosopher Theophrastus and by his Indian contemporaries **Sushrut** and **Charak**. Thus, Pteridophytes are vastly useful to humanity.

Ferns also have fertility properties. One of the famous Fern known as biofertilizeris *Azolla pinnata* R.Br. subsp. *asiatica* R.M.K. Saunders & Fowler, which shows symbiotic association with nitrogen-fixing *Anabaena azollae* Strasburger (bluegreen algae) is used extensively in the rice fields. Through this property, *Azolla*'s agronomic potential as a biofertilizer for rice has been recognized in many countries, including India. Other ferns like *Adiantum sp.*, *Asplenium sp.*, *Selaginella sp.*, *Lygodium sp.*, *Pteris sp.*, etc. are also cultivated in gardens and pots. Ferns also play a significant role in the bio-remediation of wastewater.

CONSERVATION

At present, many different ecological niches are threatened to a large extent due to a variety of anthropogenic activities, creating a serious problem for fern survival in the region. The increasing human population, expansion of agricultural land to meet the demand of food, harvesting of fuel-wood and timber, misuse of non-timber forest products (NTFPs), selective removal of species and other developmental activities are continuing at the cost of the natural flora. In addition to these, urbanization, construction of buildings, roads, housing etc. in areas, are all examples of direct attacks against nature that have resulted in the loss of pteridophytic diversity in the area. All these activities pose a threat to many epiphytic ferns, filmed ferns, climbing ferns and many other shade-loving ferns, which can result in co-extinction. The rules of the Biodiversity Convention, are powerless to address real conservation issues like this and are largely an

impediment to botanical research. The conservation of Pteridophytes has received very little attention in comparison to Angiosperms despite being medicinal, food, rituals and ornamental and ecological values. In this context, it is important to formulate and implement in situ conservation of Pteridophytes in order to maintain the richness of the biodiversity of the district and the state as a whole.

ACKNOWLEDGEMENT

We are grateful to the Joint-Director, Botanical Survey of India, Shillong and HOD, dept of Botany, Gauhati University for allowing us to access their Herbarium and Library. We are very grateful to the Karbi Anglong District Forest Officer for providing the necessary permission and facilities to explore the reserve's forest throughout the study period.

REFERENCES

- Baishya, A.K. & R.R. Rao (1982).** *Ferns and Fern-allies of Meghalaya State*, India. Scientific Publisher, Jodhpur, 162 pp.
- Barua I., Baishya A. & Neogi B. (1989).** A contribution to the pteridophytic flora of Kamrup district Assam Indian Fern J 6: 160–180 pp.
- Benniamin A. (2011).** Medicinal ferns of North-East India with special reference to Arunachal Pradesh Indian J Traditional Knowledge 10 (3): 516-522 pp.
- Bhattacharya M. K. (1994).** Certain fern flora of Karimganj District Assam New Botanist 21: 125-127 pp.
- Bhattacharya M. K., Nath a. & Baishya A. K. (1995).** Pteridophytic flora of North Cachar Hills District Assam, India Indian Fern J 12: 146-158 pp.
- Borah, P. & J. Barukial (2021).** On the pteridophytes of Bherjan-Borajan-Padumoni Wildlife Sanctuary, Assam, India. *Journal of Threatened Taxa* 13(12): 19781–19790 pp.
- Bora, P. J. & Kumar Y. (2003).** Floristic Diversity of Assam: Study of Pabitora Wildlife Sanctuary Daya Publishing House New Delhi.
- Borthakur, S.K., P. Deka & K. Nath (2001).** *Illustrated Mannual of Ferns of Assam*. Bishen Singh Mahendra Pal Singh, Dehra Dun, 468 pp.
- Borthakur, S.K., D. Baro, A. Bawri & A. Baro (2018).** *Flora of BTAD (Bodoland Territorial Area Districts, Assam), Pteridophytes and Gymnosperms*. Eastern Book House Publisher, India, Vol.1, 152 pp.
- Chandra, S. (2000).** *The Ferns of India (Enumeration, Synonyms & Distribution)*. International Book Distributors, Dehradun, India.
- Chapman, A.D. (2009).** Numbers of Living Species in Australia and the World Report for the Australian Biological Resources Study Canberra, Australia, September 2009. <http://www.environment.gov.au/biodiversity/abrs/publications/other/species/numbers/index.html>.
- Choudhury S. (2005).** Assam Flora Assam Science Technology & Environmental Council Guwahati India.
- Das, Dimbeshwar. (2020).** Diversity of pteridophytes in Nazira subdivision of Sivasagar district, Assam, N. E. India. *Indian Fern J.*29: 89-99 pp.

- Devi, J. & H.C. Majumdar (2003).** A study on Pteridophytic flora Byrnihat and Umtru area of Meghalaya State. *Plant Archives* 3(2): 207–213 pp.
- Dixit, R.D. (1984).** *A Census of the Indian Pteridophytes*. Botanical Survey of India, Howrah.
- Dutta, A.K., T.K. Dutta & K.K. Gupta (1980).** A tentative accounting of the forest flora of North Cachar Hills and Barail range-II: Enumeration of spores of Pteridophytes. *Indian Forester* 106: 34–40 pp.
- Fraser-Jenkins, C.R. (2009).** A brief comparison of modern pteridophyte classifications (families and genera in India). *Indian Fern Journal* 26: 107–126pp.
- Fraser-Jenkins, C.R., K.N. Gandhi, B.S. Kholia & A. Benniamin (2017).** *An annotated checklist of Indian pteridophytes part-1*. Bishen Singh Mahendra Pal Singh, Dehradun, India, 562 pp.
- Gangwar A. K. & Ramakrishnan P. S. (1990).** Ethno-biological Notes on Some Tribes of Arunachal Pradesh Northeastern India *Econ Bot* 44(1): 94-105 pp.
- Handique, P.J. & G. Konger (1986).** A list of Ferns and Fern allies of greater Guwahati (Assam) with their ecological adaptation. *Mendel* 3(2): 130–135 pp.
- Jain, S.K. & R.R. Rao (1977).** *A Handbook of Field and Herbarium Methods*. Today & Tomorrow's Printers and Publishers, New Delhi, 157 pp.
- Jamir, N.S. & R.R. Rao (1988).** *The Ferns of Nagaland*. Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 426 pp.
- Kachroo, P. (1953).** Ferns of Assam. *Journal of Asiatic Society, Bengal*. 29: 161–174pp.
- Kachroo, P., S.S. Bir & S.M. Vasudeva (1989).** Pteridophytic flora of North-Eastern India-II (families: Cryptogramaceae-Thelypteridaceae. *Indian Fern Journal* 6:78–99 pp.
- Kalita, P.C. (2015).** Survey for pteridophytic diversity in Madan Kamdev Hill region of Assam, India. *Pleione* 9(2): 376–400 pp.
- Maridass M.& Ghantihikumar S.(2008).** Antibacterial activity of leaves of *Blechnum orientale* L., *Pharmacology on line* 3: 58-60 pp.
- Maridass M. & Raju G. (2010).** Investigation of Phytochemical and Antimicrobial Activity of *Huperzia species* *Pharmacology on line* 3: 688-692 pp.
- Mipun, P. & Borah, Dipankar & Bhat, Nazir. (2019).** Checklist for Phytodiversity of East Karbi Anglong Wildlife Sanctuary, Assam, Northeast India: Part I. *Ambient Science*, 2019: Vol. 06(2); Online.
- Moran, R.C. (2015).** An overview of ferns and lycophytes Opening Symposium Abstract (p5) In 'Next Generation Pteridology' [An International Conference on Lycophyte and Fern Research], Smithsonian Institution and US Botanic Garden, Washington DC; June 1–5, 2015 *Book of Program and Abstracts*, 74 pp.
- Panigrahi, G. (1960).** Pteridophytes of eastern India-1 Enumeration of the species collected and their nomenclature. *Bulletin of Botanical Survey of India* 2: 309–314 pp.

Panigrahi, G. (1968). Studies in the Polypodiaceae in eastern India: distribution and ecology. *Journal of Indian Botanical Society* 47: 1–6 pp.

Panigrahi, G. & S. Choudhury (1961). Taxonomic studies on the Aspidiaceae of Eastern India. *Proceeding of Indian Science Congress* 48: 272 pp.

Panigrahi, G. & S. Choudhury (1962). Enumeration and distribution of fern allies in Eastern India. *Proceeding of Indian Science Congress* 49: 255–256 pp.

Rethy P., Singh B., Kagyung R. & Gajurel P. R. (2010). Ethno-botanical studies of Dehang – Debang Biosphere of Arunachal Pradesh with special reference to the Memba tribe Indian J Traditional Knowl 9(1): 61-67 pp.

Sen, A. & P.D. Ghosh (2011). A note of the ethnobotanical studies of some pteridophytes in Assam. *Indian Journal of Traditional Knowledge* 10(2): 292–295 pp.

Shils. & Choudhury D. M.(2009) Ethnomedicinal importance of Pteridophytes Used by the Reang tribe of Tripura North East India Ethnobot Leaflets 13: 634-643 pp.
